arXiv:math/0404147v1 [math.LO] 7 Apr 2004

LONG LOW ITERATIONS

HEIKE MILDENBERGER AND SAHARON SHELAH

Abstract. We try to control many cardinal characteristics by working with a notion of orthogonality between two families of forcings. We show that $\mathfrak{b}^+ < \mathfrak{g}$ is consistent.

This work is dedicated to James Baumgartner on the occasion of his 60th birthday.

Contents

0. Introduction 1
1. Pseudo creature forcing 3
2. Forcing bigness notions 15
3. Long low c.c.c. iteration and $\mathbf{q} \perp \Gamma$ 23
4. Examples 29
5. On the consistency of $\mathfrak{b} \ll \mathfrak{g}$ 30

References 31

0. Introduction

In this section we define cardinal characteristics that are the candidates for our arrangements. Our aim is to arrange that one characteristic be small, say be κ, and to increase the other characteristic to $\lambda > \kappa^+$. The technical means is some strategically $(< \lambda)$-complete pattern of c.c.c. forcings. If the relations underlying the two characteristics are sufficiently orthogonal, this will be possible.

For more than two characteristics at the same time, only very little is known.

Date: March 30, 2004.

2000 Mathematics Subject Classification: 03E15, 03E17, 03E35.

The first author was supported by the Austrian “Fonds zur wissenschaftlichen Förderung”, grants no. 13983 and 16334 and by travel support by the Landau Center.

The second author's research is supported by the United States-Israel Binational Science Foundation (Grant no. 2002323). This is his Publication 843.

The kinds of invariants that will be forced to be big are connected to notions of forcings $(\mathbf{Q}, \leq_{\mathbf{q}})$ that add special reals and thus increase some cardinal characteristic. Which characteristics? We give a pragmatic definition:

Definition 0.1. *Let* $\mathbf{q} = (\mathbf{Q}, \leq_{\mathbf{q}})$ *be a notion of forcing,* $\mathbf{Q} \subseteq {}^{\omega}\omega$.

(1) $\mathrm{inv}_{\mathrm{cm}}(\mathbf{q}) = \min\{\kappa$: *there is a* $\leq_{\mathbf{q}}$*-increasing sequence of length* κ *with no* $\leq_{\mathbf{q}}$ *upper bound.*$\}$.

(2) $\mathrm{inv}_{\mathrm{cm}}(\mathbf{q}, \mathrm{R}) = \min\{\kappa$: *there is a* $\leq_{\mathbf{q}}$*-increasing sequence* $\langle \bar{\mathfrak{c}}_\zeta : \zeta < \kappa\rangle$ *such that for all* $\eta \in {}^{\omega}\omega$ *there is some* $\zeta < \kappa$ *such that* $\neg \bar{\mathfrak{c}}_\zeta R \eta\}$.

(3) $\mathrm{inv}_{\mathrm{cf}}(\mathbf{q}) = \min\{|\mathcal{C}| : \mathcal{C} \subseteq \mathbf{Q} \wedge (\forall \bar{\mathfrak{d}} \in \mathbf{Q})(\exists \bar{\mathfrak{c}} \in \mathcal{C})(\bar{\mathfrak{d}} \leq_{\mathbf{q}} \bar{\mathfrak{c}})\}$.

(4) $\mathrm{inv}_{\mathrm{gm}}(\mathbf{q})$ *is the minimal* κ *such that in the following game* $\Game_\kappa(\mathbf{q})$ *the empty player has a winning strategy: A play lasts* $\kappa + 1$ *moves and in the* α*-th move the non-empty player chooses* $\bar{\mathfrak{c}}_\alpha \in \mathbf{Q}$ *satisfying* $\beta < \alpha \Rightarrow \bar{\mathfrak{d}}_\beta \leq_{\mathbf{q}} \bar{\mathfrak{c}}_\alpha$ *and then the empty player chooses* $\bar{\mathfrak{d}}_\alpha \in \mathbf{Q}$ *such that* $\bar{\mathfrak{c}}_\alpha \leq_{\mathbf{q}} \bar{\mathfrak{d}}_\alpha$. *Then the non-empty player wins iff he always has a legal move.*

Definition 0.2. *For* $\mathbf{q}$ *as in Definition 0.1 we define* $\mathrm{Spec}(\mathbf{q})$ *as the set of regular (uncountable* $\leq 2^\omega$*) cardinals* κ *such that in the game* $\Game^*_\kappa(\mathbf{q})$ *the empty player has no winning strategy. The game* $\Game^*_\kappa(\mathbf{q})$ *is defined just like* $\Game_\kappa(\mathbf{q})$ *except that in stage* κ *non-empty wins if there were* $\bar{\mathfrak{c}}_\alpha$ *for all* $\alpha < \kappa$ *but there is no* $\bar{\mathfrak{c}}_\kappa$.

Now we introduce the orthogonal relations, that shall have small characteristics: Some relations R, different from the ones in the previous definition, shall have a small R-unbounded set $\{\underset{\sim}{\eta}_i : i < \omega_1\}$ that needs to be preserved by the forcing $\mathfrak{K} * \underset{\sim}{\mathbb{P}}$, though we build a $\mathfrak{K}$-generic forcing $\underset{\sim}{\mathbb{P}}$ increasing some of the "creature"-invariants above. We work with forcing bigness notions Γ such that R fits Γ (see Definition 0.3). The $\underset{\sim}{\mathbb{P}}$ will be described by better and better approximations $(\mathbb{P}, \mathbb{P}^+, \underset{\sim}{\eta}) \in \Gamma$ on which the same $\underset{\sim}{\eta}$ keeps its rôle as a member of an R-unbounded family. Preliminarily the reader may think of a forcing bigness notion Γ as a way of finding suitable extensions in the $\lessdot$ order of the forcings. The technical definition of Γ will be given in Definition 2.1.

Definition 0.3. (1) *We say a binary Borel relation* R *on* ${}^{\omega}(\mathcal{H}(\omega))$ *fits a bigness notion* Γ *if* $(\mathbb{P}, \mathbb{P}^+, \underset{\sim}{\eta}) \in \Gamma$ *implies that*

$$\Vdash_{\mathbb{P}^+} \forall \nu \in ({}^{\omega}\omega)^{\mathbf{V}[\mathbb{P}]}(\neg \underset{\sim}{\eta} R \nu).$$

(2) *Let* $\mathrm{inv}(R) = \min\{|Y| : Y \subseteq {}^{\omega}\omega \wedge (\forall \nu \in {}^{\omega}\omega)(\exists \eta \in Y)(\neg \eta R \nu)\}$.

(3) *In* $\mathbf{V}$ *or in any generic extension* $\mathbf{V}[\mathbb{P}]$ *of* $\mathbf{V}$*, we say that* $\eta \in {}^{\omega}\omega$ *is* (Γ, R)*-big over* $A \subseteq {}^{\omega}\omega$ *if* R *is a binary Borel relation on* ${}^{\omega}(\mathcal{H}(\omega))$ *which fits* Γ *and* $\nu \in A$ *implies* $\neg \eta R \nu$*.*

(4) *Let* Γ *be a forcing bigness notion in* $\mathbf{V}_1$*,* $\mathbf{V}_2$ *extend* $\mathbf{V}_1$*,* $A \in \mathbf{V}_2$*,* $A \subseteq \mathbf{V}_1$*,* $\eta \in ({}^{\omega}\omega)^{\mathbf{V}_1}$*. We say that* η *is* Γ*-big over* A *in* $(\mathbf{V}_1, \mathbf{V}_2)$ *if for some* $(\mathbb{P}, \mathbb{P}^+, \underset{\sim}{\eta}) \in \Gamma^{\mathbf{V}_1}$ *and* $G \in \mathbf{V}_2$*,* G *is* $\mathbb{P}^+$*-generic over* $\mathbf{V}_1$*,* $A \in \mathbf{V}_1[G \cap \mathbb{P}]$*, and* $\eta = \underset{\sim}{\eta}[G]$*.*

1. Pseudo creature forcing

Now we introduce some forcings $\mathbf{q} = (\mathbf{Q}, \leq_{\mathbf{q}})$, whose invariants according to Definition 0.1 shall be increased by some compound forcings later in Section 3.

We think of forcings with linear creatures as in Shelah [13], Blass and Shelah [4], Proper and Improper Forcing [14, VI,§6] and Rosłanowski and Shelah [9].

Definition 1.1. *We say* $\mathbf{q} = (K_{\mathbf{q}}, \Sigma_{\mathbf{q}}, \mathrm{nor}_{\mathbf{q}}, \mathrm{val}_{\mathbf{q}}) = (K, \Sigma, \mathrm{nor}, \mathrm{val})$ *is a* pseudo creature forcing *(abbreviated by pcrf) iff*

(a) $K \subseteq \mathcal{H}(\aleph_0)$ *is a set of creatures of the form* $\mathfrak{c}\colon [m_{\mathrm{dn}}(\mathfrak{c}), m_{\mathrm{up}}(\mathfrak{c})) \to \mathcal{H}(\aleph_0)$*, and for each* $\mathfrak{c} \in K$ *we have* $m_{\mathrm{dn}}(\mathfrak{c}) < m_{\mathrm{up}}(\mathfrak{c}) < \omega$*.*

(b) *The subcomposition function* Σ *is a partial function from* ${}^{\omega>}K$ *to* $\mathcal{P}(K) \setminus \{\emptyset\}$ *such that*

  (α) $\mathrm{dom}(\Sigma) \subseteq \{(\mathfrak{c}_n, \dots, \mathfrak{c}_{m-1}) : n < m < \omega \wedge \forall \ell \in [n, m)(\mathfrak{c}_\ell \in K \wedge m_{\mathrm{up}}(\mathfrak{c}_\ell) \leq m_{\mathrm{dn}}(\mathfrak{c}_{\ell+1}))\}$.

  (β) $\Sigma(\mathfrak{c}_n, \dots, \mathfrak{c}_{m-1}) \subseteq \{\mathfrak{c} \in K : (\exists k, \ell)(n \leq k < \ell \leq m \wedge m_{\mathrm{dn}}(\mathfrak{c}) = m_{\mathrm{dn}}(\mathfrak{c}_k), m_{\mathrm{up}}(\mathfrak{c}) = m_{\mathrm{up}}(\mathfrak{c}_{\ell-1}))\}$.

  (γ) Σ *is transitive: If* $\mathfrak{c}^1_\ell \in \Sigma(\mathfrak{c}^0_{m_\ell}, \dots, \mathfrak{c}^0_{m_{\ell+1}-1})$ *for* $\ell < k$ *and* $\mathfrak{c}^2 \in \Sigma(\mathfrak{c}^1_0, \dots, \mathfrak{c}^1_{k-1})$*, then* $\mathfrak{c}^2 \in \Sigma(\mathfrak{c}^0_{m_0}, \dots, \mathfrak{c}^0_{m_1-1}, \mathfrak{c}^1_{m_1}, \dots, \mathfrak{c}^1_{m_2-1}, \dots, \mathfrak{c}^{k-1}_{m_k-1})$*. That also means, that the domain of* Σ *is closed under the mentioned concatenations.*

  (δ) $(\forall \mathfrak{c} \in K)(\mathfrak{c} \in \Sigma(\mathfrak{c}))$ *and* Σ *is monotonous in the following sense: If* $i_0 < \dots < i_{m-1} < n$ *then*

$$\Sigma(\mathfrak{c}_{i_0}, \dots, \mathfrak{c}_{i_{n-1}}) \subseteq \Sigma(\mathfrak{c}_0, \dots, \mathfrak{c}_{n-1}).$$

(c) $\mathrm{nor}\colon K \to \mathbb{R}^{\geq 0}$ *(usually* ω*).*

(d) $\mathrm{val}\colon K \to \omega^{<\aleph_0}$*,* $\emptyset \neq \mathrm{val}(\mathfrak{c}) \subseteq [m_{\mathrm{dn}}(\mathfrak{c}), m_{\mathrm{up}}(\mathfrak{c}))$*.*

Remark: Note that Definition 1.1. (b)(β) is a stronger requirement on the subcomposition functions than the usual $m_{\rm dn}(\mathfrak{c}) \geq m_{\rm dn}(\mathfrak{c}_n)$, $m_{\rm up}(\mathfrak{c}) \leq m_{\rm up}(\mathfrak{c}_{m-1})$.

Definition 1.2. *1. The set of pure conditions of* $\mathbf{q}$ *is*

$$\begin{aligned} \mathbf{Q}^{pr}_{\mathbf{q}} = \{\bar{\mathfrak{c}} \,:\, \bar{\mathfrak{c}} =& \langle \mathfrak{c}_n \,:\, n<\omega\rangle, \mathfrak{c}_n \in K, m_{\rm up}(\mathfrak{c}_n) = m_{\rm dn}(\mathfrak{c}_{n+1}), m_{\rm dn}(\mathfrak{c}_0) = 0 \\ & \wedge \lim_{n\to\omega} {\rm nor}(\mathfrak{c}_n) = \infty\}. \end{aligned} \tag{1.1}$$

We usually write $\bar{\mathfrak{c}} \in K$. *Let* $\mathbf{Q}^p_{\mathbf{q}}$ *be defined similarly, allowing* $m_{\rm dn}(\mathfrak{c}_0) > 0$.

2. The set of not necessarily pure conditions $\mathbf{Q}^{np}_{\mathbf{q}}$ *of* $\mathbf{q}$ *is*

$$\begin{aligned} \mathbf{Q}^{np}_{\mathbf{q}} = \{\bar{\mathfrak{c}} \,:\, \bar{\mathfrak{c}} =& \langle \mathfrak{c}_n \,:\, n<\omega\rangle, \mathfrak{c}_n \in K, m_{\rm up}(\mathfrak{c}_n) \leq m_{\rm dn}(\mathfrak{c}_{n+1}), \\ & \wedge \lim_{n\to\omega} {\rm nor}(\mathfrak{c}_n) = \infty\}. \end{aligned} \tag{1.2}$$

3. The true creature forcing $\mathbf{Q}^{tr}_{\mathbf{q}}$ *is a subset of*

$$\begin{aligned} \{\langle w, \mathfrak{c}_0, \mathfrak{c}_1, \dots \rangle \,:\, & (\exists n_0 < n_1 < \dots)(\forall i<\omega)(\exists u_i \subseteq [n_i, n_{i+1}))(\exists \bar{\mathfrak{c}}^* \in \mathbf{Q}^{np}_{\mathbf{q}}) \\ & (u_i \neq \emptyset \wedge w \in {\rm pos}(\langle\rangle, \mathfrak{c}^*_0, \dots \mathfrak{c}^*_{n_0-1}) \wedge \mathfrak{c}_i \in \Sigma(\mathfrak{c}^*_m)_{m\in u_i})\}, \end{aligned}$$

This is not quite the general case of creature forcing. In the terminology of [9, Definition 2.1.1.] *all our forcings are omittory. The pos (possibility) operation is defined as follows:*

$$\begin{aligned} {\rm pos}(w, \mathfrak{c}_0, \dots \mathfrak{c}_{n-1}) =& \{ w \cup \bigcup_{\ell<k} w_\ell \,:\, \exists \langle u_\ell, \mathfrak{d}_\ell \,:\, \ell < k\rangle, u_\ell \neq \emptyset, u_\ell \subseteq n, \\ & \max(u_\ell) < \min(u_{\ell+1}), \mathfrak{d}_\ell \in \Sigma((\mathfrak{c}_i)_{i\in u_\ell}), w_\ell \in {\rm val}(\mathfrak{d}_\ell)\}. \end{aligned} \tag{1.3}$$

Remark 1.3. $\mathbf{Q}^{pr}_{\mathbf{q}}$ and $\mathbf{Q}^p_{\mathbf{q}}$ are the pure elements in [9]. Pure means: $m_{\rm up}(\bar{\mathfrak{c}}_n) = m_{\rm dn}(\bar{\mathfrak{c}}_{n+1})$ for $n<\omega$, so the union of the domains is ω in the case of $\mathbf{Q}^{pr}_{\mathbf{q}}$ and $\omega \setminus n$ for some n in the case of $\mathbf{Q}^p_{\mathbf{q}}$. Of course, one could also introduce a pure true creature forcing. But for our intended applications, pureness is often too strong a requirement.

Now we equip $\mathbf{Q}^{pr}_{\mathbf{q}}$ and $\mathbf{Q}^p_{\mathbf{q}}$ with two partial orderings $\leq_{\mathbf{q}}$ and the stricter $\leq_{\rm full}$.

Definition 1.4. *1. For* $\bar{\mathfrak{c}}^1, \bar{\mathfrak{c}}^2 \in \mathbf{Q}^{pr}_{\mathbf{q}}$ *or in* $\mathbf{Q}^p_{\mathbf{q}}$ *we write* $\bar{\mathfrak{c}}^1 \leq^{pr}_{\mathbf{q}} \bar{\mathfrak{c}}^2$ *iff there is some* $i(*) \in \omega$ *and there is some sequence* $n_{i(*)} < n_{i(*)+1} < \dots$ *such that for every* $i \geq i(*)$ $\mathfrak{c}^2_i \in \Sigma(\mathfrak{c}^1_{n_i}, \dots, \mathfrak{c}^1_{n_{i+1}-1})$ *and* $m_{\rm dn}(\mathfrak{c}^2_i) = m_{\rm dn}(\mathfrak{c}^1_{n_i})$ *and* $m_{\rm up}(\mathfrak{c}^2_i) = m_{\rm up}(\mathfrak{c}^1_{n_{i+1}-1})$ *(so, and in Definition 1.1(b)(β) there is equality in the pure case).*

We say that $(i(*), n_{i(*)})$ *witnesses* $\bar{\mathfrak{c}}^1 \leq^{pr}_{\mathbf{q}} \bar{\mathfrak{c}}^2$.

If we have in addition $i(*)=0$ *(then, in case* $\bar{\mathfrak{c}}^1,\bar{\mathfrak{c}}^2\in\mathbf{Q}^{\rm pr}_{\mathbf{q}}$*, also* $n_{i(*)}=0$*), then we write* $\bar{\mathfrak{c}}^1\leq^{pr}_{\rm full}\bar{\mathfrak{c}}^2$*. The reversed relations are* $\geq^{pr}_{\mathbf{q}}$ *and* $\geq^{pr}_{\rm full}$ *respectively.*

2. For $\bar{\mathfrak{c}}^1,\bar{\mathfrak{c}}^2\in\mathbf{Q}^{np}_{\mathbf{q}}$ *we write* $\bar{\mathfrak{c}}^1\leq^{np}_{\mathbf{q}}\bar{\mathfrak{c}}^2$ *iff there is some* $i(*)\in\omega$ *and there is some sequence* $n_{i(*)}<n_{i(*)+1}<\dots$ *and some sequence* $\emptyset\neq u_i\subseteq[n_i,n_{i+1})$ *such that for every* $i\geq i(*)$*,* $\mathfrak{c}^2_i\in\Sigma((\mathfrak{c}^1_z)_{z\in u_i})$*. So* $m_{\rm dn}(\mathfrak{c}^2_i)=m_{\rm dn}(\mathfrak{c}^1_{\min(u_i)})$*,* $m_{\rm up}(\mathfrak{c}^2_i)=m_{\rm dn}(\mathfrak{c}^1_{\max(u_i)})$*.*

3. For $(w^1,\bar{\mathfrak{c}}^1),(w^2,\bar{\mathfrak{c}}^2)\in\mathbf{Q}^{tr}_{\mathbf{q}}$ *we write* $(w^1,\bar{\mathfrak{c}}^1)\leq^{tr}_{\mathbf{q}}(w^2,\bar{\mathfrak{c}}^2)$ *iff there is some some sequence* $0\leq n_0<n_1<\dots$ *and some sequence* $u_i\subseteq[n_i,n_{i+1})$ *such that for every* $i\in\omega$*,* $\mathfrak{c}^2_i\in\Sigma((\mathfrak{c}^1_z)_{z\in u_i})$*, and such that* $w^1\trianglelefteq w^2$ *and* $w^2\setminus w^1\in{\rm pos}(\bar{\mathfrak{c}}^1\restriction n_0)$*.*

Convention 1.5. *1. In the pure context, we write* $\langle\mathfrak{c}_{n_0},\dots,\mathfrak{c}_{n_1-1}\rangle\in\Sigma(\mathfrak{c}_{k_0},\dots,\mathfrak{c}_{k_1-1})$ *if there are* $k_0=m_0<m_1<\dots<m_{n_1}=k_1$ *such that for all* $j\in[n_0,n_1)$*,* $\mathfrak{c}_j\in\Sigma(\mathfrak{c}_{m_j},\dots,\mathfrak{c}_{m_{j+1}-1})$*.*

2. In the not necessarily pure context, we write $\langle\mathfrak{c}_{n_0},\dots,\mathfrak{c}_{n_1-1}\rangle\in\Sigma(\mathfrak{c}_{k_0},\dots,\mathfrak{c}_{k_1-1})$ *if there are* $k_0=m_0<m_1<\dots<m_{n_1}=k_1$ *and* $\emptyset\neq u_i\subseteq[m_i,n_{i+1})$ *such that for all* $j\in[n_0,n_1)$*,* $\mathfrak{c}_j\in\Sigma((\mathfrak{c}_z)_{z\in u_i})$*.*

The following property can be used for all the variants of notions of forcing $(\mathbf{Q}^{xy}_{\mathbf{q}},\leq^{xy}_{\mathbf{q},{\rm full}})$ so far defined.

Definition 1.6. *We say that a pcrf* $\mathbf{q}$ *is forgetful if the following holds: If* $\bar{\mathfrak{c}}^1\leq_{\mathbf{q}}\bar{\mathfrak{c}}^2$ *and* $n\leq k<\omega$ *and* $\bar{\mathfrak{c}}^1\restriction k=\bar{\mathfrak{c}}^2\restriction k$ *and* $\bar{\mathfrak{e}}^\ell\in\Sigma(\bar{\mathfrak{c}}^1\restriction[n,k))$ *for* $\ell=1,2$*, then we can find* m_1,m_2 *such that* $m_{\rm dn}(\mathfrak{c}^1_{m_1})=m_{\rm dn}(\mathfrak{c}^2_{m_2})$*,* $m_1>k$*,* $m_2>n$ *and*

$$\bigcap_{\ell=1}^{2}\Sigma(\bar{\mathfrak{e}}^{\ell\frown}\bar{\mathfrak{c}}^1\restriction[k,m_1))\cap\Sigma(\bar{\mathfrak{c}}^2\restriction[n,m_2))\neq\emptyset.$$

Note that ${\rm lg}(\bar{\mathfrak{e}}^\ell)\leq k-n$ and may be strictly less, and that

$$(\bar{\mathfrak{e}}^{\ell\frown}\bar{\mathfrak{c}}^1)(i)=\begin{cases}\mathfrak{e}^\ell_i, & \text{if } i<{\rm lg}(\bar{\mathfrak{e}}^\ell),\\ \mathfrak{c}^1_{k+j}, & \text{if } i={\rm lg}(\bar{\mathfrak{e}}^\ell)+j.\end{cases}$$

but since we do not need to look so close at the conditions this shift in indexing will not appear explicitly.

Convention 1.7. *We let* $\mathbf{q}$ *denote a forgetful pcrf unless said otherwise.*

Definition 1.8. *We define an equivalence relation* $\equiv_{\mathbf{q}}$ *on* $\mathbf{Q}^{np}_{\mathbf{q}}$ *and on* $\mathbf{Q}^{tr}_{\mathbf{q}}$*:* $\bar{\mathfrak{c}}^1\equiv_{\mathbf{q}}\bar{\mathfrak{c}}^2$ *iff there are* $k_1,k_2\in\omega$ *such that for all* $n<\omega$*,* $\mathfrak{c}^1_{k_1+n}=\mathfrak{c}^2_{k_2+n}$*.*

All equivalence classes of this relation are countable. In the language of Kechris, Hjorth, Louveau and others this is a countable equivalence relation.

Definition 1.9. *Let* $\mathbf{q} = (\mathbf{Q}_{\mathbf{q}}, \leq_{\mathbf{q}})$ *any of our versions of forcings.*

1. $\mathcal{C}$ *is a* $\mathbf{q}$*-pre-directed family if*
   - (a) $\mathcal{C} \subseteq \mathbf{Q}_{\mathbf{q}}$,
   - (b) *for some proper filter* D *on* ω *containing the cofinite sets of* ω*, any finitely many members of* $\mathcal{C}$ *have a common upper bound* $\bar{\mathfrak{c}}$ *w.r.t.* $\leq_{\mathbf{q}}$ *with* $\mathrm{supp}(\bar{\mathfrak{c}}) = \{m_{\mathrm{dn}}(\mathfrak{c}_n) \,:\, n \in \omega\} \in D$.

   Note that in (b) we did not require that $\mathfrak{c} \in \mathcal{C}$*. This explains the prefix "pre". Such a filter* D *is called a witness for* $\mathcal{C}$ *being* $\mathbf{q}$*-pre-directed.*

2. $\mathcal{C}$ *is a* $\mathbf{q}$*-directed family if*
   - (a) $\mathcal{C} \subseteq \mathbf{Q}_{\mathbf{q}}$,
   - (b) $(\mathcal{C}, \leq_{\mathbf{q}})$ *is directed, i.e., every two members of* $\mathcal{C}$ *have a common upper bound in* $\mathcal{C}$ *w.r.t.* $\leq_{\mathbf{q}}$,
   - (c) *If* $\bar{\mathfrak{c}}^1 \equiv_{\mathbf{q}} \bar{\mathfrak{c}}^2 \in \mathbf{Q}_{\mathbf{q}}$*, then* $\bar{\mathfrak{c}}^1 \in \mathcal{C}$ *iff* $\bar{\mathfrak{c}}^2 \in \mathcal{C}$.

In the following lemma we use the strong requirement from Definition 1.1(b)(β).

Lemma 1.10. *Let the order be* $\leq^{pr}_{\mathbf{q}}$, $\leq^{np}_{\mathbf{q}}$*, or* $\leq^{tr}_{\mathbf{q}}$*. Every* $\mathbf{q}$*-directed family is* $\mathbf{q}$*-pre-directed.*

Proof. Let $\mathcal{C}$ be $\mathbf{q}$-directed. Take D be the filter generated by $\{\mathrm{supp}(\bar{\mathfrak{c}}) \,:\, \bar{\mathfrak{c}} \in \mathcal{C}\}$. If $\bar{\mathfrak{c}}^1 \leq^{pr}_{\mathbf{q}} \bar{\mathfrak{c}}^2$, then $\mathrm{supp}(\bar{\mathfrak{c}}^2) \subseteq^* \mathrm{supp}(\bar{\mathfrak{c}}^1)$, and for $\leq^{pr}_{\mathrm{full}}$ there are no exceptions to the inclusion. □

Definition 1.11. *Note that* $\mathbb{A}$ *is not identical with* $\mathbf{Q}$.

For a $\mathbf{q}$*-directed family* $\mathcal{C}$ *in* $(\mathbf{Q}_{\mathbf{q}}, \leq_{\mathbf{q}})$ *we define a partial order* $\mathbb{A}_{\mathbf{q}}(\mathcal{C}) = \mathbb{A}(\mathcal{C})$ *as follows:*

- (a) *the elements are pairs* $(n, \bar{\mathfrak{c}})$ *where* $n < \omega$ *and* $\bar{\mathfrak{c}} \in \mathcal{C}$,
- (b) *the order is* $(n_1, \bar{\mathfrak{c}}^1) \leq_{\mathbb{A}(\mathcal{C})} (n_2, \bar{\mathfrak{c}}^2)$ *iff*
  - (α) $n_1 \leq n_2$,
  - (β) $\bar{\mathfrak{c}}^1 \restriction n_1 = \bar{\mathfrak{c}}^2 \restriction n_1$,

- *(γ)* $\bar{\mathfrak{c}}^1 \restriction [n_1, \omega) \leq_{\text{full}} \bar{\mathfrak{c}}^2 \restriction [n_1, \omega)$, *which means, by the definition of* $\leq_{\text{full}}$*: for some* $m_0 = n_1 < m_1 < m_2 \ldots$ *we have for all* $i < \omega$, $\mathfrak{c}^2_{n_1+i} \in \Sigma(\mathfrak{c}^1_{m_i}, \ldots, \mathfrak{c}^1_{m_{i+1}-1})$.

Often we write only $\leq$ *for* $\leq_{\mathbb{A}(\mathcal{C})}$.

Definition 1.12. *Let* $\mathcal{C}$ *be* $\mathbf{q}$*-directed. For* $\mathfrak{c} \in \mathcal{C}$*, let* $\mathbf{Q}(\bar{\mathfrak{c}}) = \{\bar{\mathfrak{c}}' \in \mathbf{Q} : \bar{\mathfrak{c}}' \equiv_{\mathbf{q}} \bar{\mathfrak{c}}\}$ *and let* $\mathbb{A}(\bar{\mathfrak{c}}) = \{(n, \bar{\mathfrak{c}}') : n < \omega, \bar{\mathfrak{c}}' \in \mathbf{Q}(\bar{\mathfrak{c}})\}$*. Let* $\mathbf{Q}_{\mathcal{C}}(\bar{\mathfrak{c}}) = \{\bar{\mathfrak{c}}' \in \mathcal{C} : \bar{\mathfrak{c}}' \equiv_{\mathbf{q}} \bar{\mathfrak{c}}\}$ *and let* $\mathbb{A}_{\mathcal{C}}(\bar{\mathfrak{c}}) = \{(n, \bar{\mathfrak{c}}') : n < \omega, \bar{\mathfrak{c}}' \in \mathbf{Q}_{\mathcal{C}}(\bar{\mathfrak{c}})\}$*. Each* $\equiv_{\mathbf{q}}$*-class is infinite and countable. Hence, if above each condition there are two incompatible conditions, then the second and the fourth forcing (with* $\leq_{\mathbb{A}(\mathcal{C})}$*) are equivalent to the Cohen forcing.*

Definition 1.13. *We say that* $\mathcal{C}$ *is strongly* $\mathbf{q}$*-directed if* $(\mathcal{C}, \leq_{\mathbf{q}})$ *is* $\mathbf{q}$*-directed and* $\aleph_1$*-directed, that is, every countable set has a common upper bound.*

Claim 1.14. *Let* $\mathbf{q}$ *be a forgetful pcrf.*

1. *If* $\mathcal{C}$ *is* $\mathbf{q}$*-directed, then* $\mathbb{A}(\mathcal{C})$ *is a* σ*-centered forcing notion.*
2. *If* $\mathcal{C}$ *is* $\mathbf{q}$*-pre-directed and* $\bar{\mathfrak{c}}^n \in \mathcal{C}$ *for* $n < \omega$ *(possibly with repetitions), then*

$$\Vdash_{\text{Cohen}} \text{“for some } \bar{\mathfrak{c}} \in \mathbf{Q_q}^{\mathbf{V}[\text{Cohen}]} \text{ we have } n < \omega \rightarrow \bar{\mathfrak{c}}^n \leq_{\mathbf{q}} \bar{\mathfrak{c}}$$
$$\text{and } \mathcal{C} \cup \{\bar{\mathfrak{c}}\} \text{ is } \mathbf{q}\text{-pre-directed.”}$$

3. Let $\mathcal{C}$ *be a* $\mathbf{q}$*-pre-directed family and* $\mathbb{P}$ *be the forcing of adding* $2^{\aleph_0}$ *Cohen reals. Then in* $\mathbf{V}^{\mathbb{P}}$ *we can extend* $\mathcal{C}$ *to a strongly* $\mathbf{q}$*-directed* $\mathcal{C}'$.

4. If $\mathcal{C}$ *is strongly* $\mathbf{q}$*-directed and* $\mathcal{I}$ *is a predense subset of* $\mathbb{A}(\mathcal{C})$*, then for some* $\bar{\mathfrak{c}} \in \mathcal{C}$ *and some* $\mathcal{J}$ *we have*

- *(α)* $\mathcal{J} \subseteq \mathbb{A}_{\mathcal{C}}(\bar{\mathfrak{c}})$*, hence is countable,*
- *(β)* $\mathcal{J}$ *is predense in* $\mathbb{A}(\mathcal{C})$*,*
- *(γ)* $(\forall q \in \mathcal{J})(\exists p \in \mathcal{I})(p \leq_{\mathbb{A}(\mathcal{C})} q)$.

5. If $\mathcal{C}^1 \subseteq \mathcal{C}^2$ *are* $\mathbf{q}$*-directed, then*

- *(α) If* $p, q \in \mathbb{A}(\mathcal{C}^1)$ *and* $\mathbb{A}(\mathcal{C}^1) \models p \leq q$ *then* $\mathbb{A}(\mathcal{C}^2) \models p \leq q$*,*
- *(β) If* $p, q \in \mathbb{A}(\mathcal{C}^1)$ *and* $\mathbb{A}(\mathcal{C}^1) \models p \not\leq q$ *then* $\mathbb{A}(\mathcal{C}^2) \models p \not\leq q$*,*
- *(γ) If* $p = (n^p, \bar{\mathfrak{c}}^p)$*,* $q = (n^q, \bar{\mathfrak{c}}^q) \in \mathbb{A}(\mathcal{C}^1)$ *are incompatible in* $\mathbb{A}(\mathcal{C}^1)$ *and* $\bar{\mathfrak{c}}^p \leq_{\mathbf{q}} \bar{\mathfrak{c}}^q$ *then* p, q *are incompatible in* $\mathbb{A}(\mathcal{C}^2)$.
- *(δ) If* $p = (n^p, \bar{\mathfrak{c}}^p)$*,* $q = (n^q, \bar{\mathfrak{c}}^q) \in \mathbb{A}(\mathcal{C}^1)$ *are incompatible in* $\mathbb{A}(\mathcal{C}^1)$*, then* p, q *are incompatible in* $\mathbb{A}(\mathcal{C}^2)$.

6. *The family of* $\mathbf{q}$*-pre-directed families* $\mathcal{C}$ *as well as the family of* $\mathbf{q}$*-directed families* $\mathcal{C}$ *are closed under increasing unions.*

Proof. 1. Choose $\bar{\mathfrak{c}}^* \in \mathcal{C}$ and fix it for the rest of the proof. (Since $\mathcal{C}$ is $\mathbf{q}$-directed, the choice does not matter.)

We set

$$Y = \{(\bar{\mathfrak{d}}, n_0, n_1, n_2) \,:\, n_0 \leq n_1 < \omega, n_2 \in \omega, \bar{\mathfrak{d}} \in {}^{n_1}K, \}.$$

For each $y = (\bar{\mathfrak{d}}, n_0, n_1, n_2) = (\bar{\mathfrak{d}}^y, n_0^y, n_1^y, n_2^y) \in Y$ we define

$$\begin{aligned} \mathbb{A}_y^0 = \{p \in \mathbb{Q}(\mathcal{C}) \;: n^p = n_0, \bar{\mathfrak{c}}^p \restriction n_1 = \bar{\mathfrak{d}}, \bar{\mathfrak{c}}^* \restriction [n_2, \omega) \leq_{\text{full}} \bar{\mathfrak{c}}^p \restriction [n_1, \omega), \\ \text{so, in particular } m_{\text{dn}}(\mathfrak{c}^*_{n_2}) = m_{\text{dn}}(\mathfrak{c}^p_{n_1})\} \end{aligned} \tag{1.4}$$

and

$$\mathbb{A}_y = \{p \in \mathbb{A}(\mathcal{C}) \,:\, (\exists q \in \mathbb{A}_y^0)(p \leq q)\}. \tag{1.5}$$

Now, since $\bar{\mathfrak{d}} \in {}^{\omega>}K$ and $K \subseteq \mathcal{H}(\omega)$ is countable, we have that Y is countable.

Now we prove that $\mathbb{A}(\mathcal{C}) = \bigcup\{\mathbb{A}_y \,:\, y \in Y\}$. Let $p \in \mathbb{A}(\mathcal{C})$. Recalling that $\mathcal{C}$ is $\leq_{\mathbf{q}}$-directed, there is some $\bar{\mathfrak{c}}^1 \in \mathcal{C}$ such that $\bar{\mathfrak{c}}^p \leq_{\mathbf{q}} \bar{\mathfrak{c}}^1$ and $\bar{\mathfrak{c}}^* \leq_{\mathbf{q}} \bar{\mathfrak{c}}^1$. By the definition of $\leq_{\mathbf{q}}$ and considering, that once $i(*)$ is large enough, all larger i could serve as well, we find k_0, k_1, k_2 such that (k_0, k_2) witness $\bar{\mathfrak{c}}^p \leq_{\mathbf{q}} \bar{\mathfrak{c}}^1$ and $k_0 \leq k_1 = n_{i(*)}$ and (k_1, k_2) witness $\bar{\mathfrak{c}}^* \leq_{\mathbf{q}} \bar{\mathfrak{c}}^1$. We choose $k_0 \geq n^p$.

Now let $y = (\bar{\mathfrak{c}}^p \restriction k_1, k_0, k_1, k_2)$. Easily $y \in Y$, and let $\bar{\mathfrak{c}}^2 = (\bar{\mathfrak{c}}^p \restriction k_1, \mathfrak{c}^1_{k_2}, \mathfrak{c}^1_{k_2+1}, \dots)$ and $p_2 = (k_0, \bar{\mathfrak{c}}^2)$. Now, since $\mathfrak{c} \in \Sigma(\mathfrak{c})$ and since $k_0 \geq n^p$ and $k_0 \geq n_{i(*)}$ for $\bar{\mathfrak{c}}^p \leq_{\mathbf{q}} \bar{\mathfrak{c}}^1$, we have $\bar{\mathfrak{c}}^p \restriction k_0 = \bar{\mathfrak{c}}^2 \restriction k_0$ and $\bar{\mathfrak{c}}^p \restriction [k_0, \infty) \leq \bar{\mathfrak{c}}^2 \restriction [k_1, \infty)$, hence $p \leq p_2 \in \mathbb{A}(\mathcal{C})$, since $\bar{\mathfrak{c}}^1 \restriction [k_2, \infty) \leq_{\text{full}} \bar{\mathfrak{c}}^2[k_1, \infty)$ we have $p_2 \in \mathbb{A}_y^0$, hence $p \in \mathbb{A}_y$ as required.

Each $\mathbb{A}_y$ is directed in $\leq = \leq_{\mathbb{A}(\mathcal{C})}$: Let $y = (\bar{\mathfrak{d}}, n_0, n_1, n_2)$ and let $p_1, p_2 \in \mathbb{A}_y$. We can find $q_1, q_2 \in \mathbb{A}_y^0$ such that $p_\ell \leq q_\ell$ for $\ell = 1, 2$. So $\bar{\mathfrak{c}}^{q_1}, \bar{\mathfrak{c}}^{q_2} \in \mathcal{C}$ and we can find $\bar{\mathfrak{c}} \in \mathcal{C}$ such that $\bar{\mathfrak{c}}^{q_\ell} \leq_{\mathbf{q}} \bar{\mathfrak{c}}$ for $\ell = 1, 2$. Hence there are $k_1, k_2, k \in \omega$ such that (k_ℓ, k) witnesses $\bar{\mathfrak{c}}^{q_\ell} \leq_{\mathbf{q}} \bar{\mathfrak{c}}$ for $\ell = 1, 2$ and $k \geq n^{q_\ell}$, $k_\ell \geq n_2$. Since $q_\ell \in \mathbb{A}_y^0$, for some $k^* > n_1$ we have

$$\bar{\mathfrak{c}}^{q_\ell} \restriction [n_1, k_\ell) \in \Sigma(\bar{\mathfrak{c}}^* \restriction [n_2, k^*)) \text{ for } \ell = 1, 2.$$

Without loss of generality $\bar{\mathfrak{c}} \restriction n_1 = \bar{\mathfrak{c}}^* \restriction n_1$. Recall $\bar{\mathfrak{c}}^{q_\ell} \restriction n_1 = \bar{\mathfrak{d}}$. Now apply the definition of forgetfulness with $n_2, n_1, \bar{\mathfrak{c}}^{q_\ell} \restriction [n_1, k_\ell)(\ell = 1, 2), \bar{\mathfrak{c}}^*, \bar{\mathfrak{c}}$ standing for $n, k, \bar{\mathfrak{e}}_\ell(\ell = 1, 2), \bar{\mathfrak{c}}^1, \bar{\mathfrak{c}}^2$ gives m_1 and m_2 and $\bar{\mathfrak{c}}'$ in the intersection. Let $\bar{\mathfrak{c}}'' = (\bar{\mathfrak{d}}{}^\frown\bar{\mathfrak{c}}'{}^\frown\bar{\mathfrak{c}} \restriction [m_2, \omega))$. Then $(n_0, \bar{\mathfrak{c}}'') \in \mathbb{A}_y^0$ is a common upper bound of $q_1, q_2 \in \mathbb{A}_y^0$ in

$\leq$. Since the upper bound is in $\mathbb{A}_y$, we can iterate the process and find common upper bounds in $\mathbb{A}_y$ for finitely many $p_1, \dots, p_n$.

2. We shall work on the following two cases

(A) $\{\bar{\mathfrak{c}}^n : n \in \omega\}$ has just two members $\bar{\mathfrak{c}}^0$ and $\bar{\mathfrak{c}}^1$.

(B) For all n, $\bar{\mathfrak{c}}^n \leq_{\mathbf{q}} \bar{\mathfrak{c}}^{n+1}$.

In both cases we define a countable notion of forcing $\mathbb{P}$ that adds a solution. First we show that to consider (A) and (B) is enough. We set $\bar{\mathfrak{c}}^0 = \bar{\mathfrak{c}}^0$ and, for $n > 0$ we iteratively add $\bar{\mathfrak{d}}^n \geq_{\mathbf{q}} \bar{\mathfrak{c}}^n, \bar{\mathfrak{d}}^{n-1}$. Thereafter we add $\bar{\mathfrak{d}}^\omega$ above all $\bar{\mathfrak{d}}^n$'s. So we add $\omega + 1$ times a Cohen real and perform this iteration with finite supports, and this is again equivalent to Cohen forcing. Again forgetfulness makes it work.

In case (A) we choose $m^\ell_0 < m^\ell_1 < \dots$ such that $m_{\mathrm{dn}}(\mathfrak{c}^0_{m^0_i}) = m_{\mathrm{dn}}(\mathfrak{c}^1_{m^1_i})$ for $i < \omega$ and such that $\{m_{\mathrm{dn}}(\mathfrak{c}^0_{m^0_i}) : i < \omega\} \in D$. Now, by forgetfulness, the Cohen forcing, in its disguised form $(\{\langle \mathfrak{d}_0, \dots \mathfrak{d}_{k-1}\rangle : (\forall \ell < k)\mathfrak{d}_\ell \in \Sigma(\bar{\mathfrak{c}}^0 \restriction [m^0_\ell, m^0_{\ell+1})) \cap \Sigma(\bar{\mathfrak{c}}^1 \restriction [m^1_\ell, m^1_{\ell+1}))\}, \trianglelefteq)$ will add a generic $\bar{\mathfrak{d}}$ such that $\mathrm{supp}(\bar{\mathfrak{d}}) \in D$ and such that $\bar{\mathfrak{d}} \geq_{\mathbf{q}} \bar{\mathfrak{c}}^0, \bar{\mathfrak{c}}^1$.

In case (B) we force with Cohen forcing in the guise $(\bigcup_{n<\omega} \mathbb{A}_{\mathcal{C}}(\bar{\mathfrak{d}}^n), \leq_{\mathbb{A}(\mathcal{C})})$. By density arguments, the generic $\bar{\mathfrak{d}}$ will fulfill $\bar{\mathfrak{d}} \geq_{\mathbf{q}} \bar{\mathfrak{d}}^n$ for all n. Note that we did not enlarge the filter D.

3. We repeat part 2. $2^{\aleph_0}$ times, by bookkeeping adding common upper bounds to all countable subsets of all intermediate $\mathcal{C}_1 \supseteq \mathcal{C}$ appearing along the iteration and such that only $\mathbf{q}$-pre-directed families appear along the iteration.

4. Let $\mathcal{J}^+ = \{(n, \bar{\mathfrak{c}}) \in \mathbb{A}(\mathcal{C}) : (n, \bar{\mathfrak{c}})$ is above some member of $\mathcal{I}\}$. This is dense and open, and let $\mathcal{J}'$ be a maximal set of pairwise incompatible elements of $\mathcal{J}^+$. By part 1. of this claim, $\mathcal{J}'$ is countable. Hence $\{\bar{\mathfrak{c}}^p : p \in \mathcal{J}'\}$ is countable. Since $\mathcal{C}$ is strongly $\mathbf{q}$-directed, there is a $\leq_{\mathbf{q}}$-upper bound $\bar{\mathfrak{c}}^*$ of $\{\bar{\mathfrak{c}}^p : p \in \mathcal{J}'\}$. Finally we set

$$\mathcal{J} = \{p' \in \mathbb{A}_{\mathcal{C}}(\bar{\mathfrak{c}}^*) : p' \text{ is above some member of } \mathcal{J}'\}.$$

Clearly $\mathcal{J}$ satisfies requirements (α) and (γ). As for (β), let $r \in \mathbb{A}(\mathcal{C})$ so r is compatible with some $p \in \mathcal{J}'$. Choose $r^+ \in \mathbb{A}(\mathcal{C})$ such that $r^+ \geq r, p$. Since $\bar{\mathfrak{c}}^{r^+}$ and $\bar{\mathfrak{c}}^*$ are compatible in $(\mathcal{C}, \leq_{\mathbf{q}})$, there is a common upper bound $\bar{\mathfrak{d}} \in \mathcal{C}$. From the definitions of $\bar{\mathfrak{c}}^p \leq^* \bar{\mathfrak{c}}^{r^+} \leq_{\mathbf{q}} \bar{\mathfrak{d}}$ and of $\bar{\mathfrak{c}}^p \leq_{\mathbf{q}} \bar{\mathfrak{c}}^*$ we get series $\langle m^\ell_i : i < \omega\rangle$,

$\ell = 0, 1, 2, 3$ such that for every i,

$$\bar{\mathfrak{c}}^* \restriction [m^1_i, m^1_{i+1}) \in \Sigma(\bar{\mathfrak{c}}^p \restriction [m^0_i, m^0_{i+1})) \text{ and}$$
$$\bar{\mathfrak{d}} \restriction [m^3_i, m^3_{i+1}) \in \Sigma(\bar{\mathfrak{c}}^* \restriction [m^1_i, m^1_{i+1})) \cap \Sigma(\bar{\mathfrak{c}}^{r^+} \restriction [m^2_i, m^2_{i+1})).$$

W.l.o.g., $m^2_0 \geq n^{r^+}$ and $m^2_0 \geq n^p$. Then $q = (m^2_0, (\bar{\mathfrak{c}}^{r^+} \restriction m^2_0){}^\frown \bar{\mathfrak{d}} \restriction [m^3_0, \omega))$. Now $q \geq p' = (n^p, \bar{\mathfrak{c}}^p \restriction n^p{}^\frown \bar{\mathfrak{c}}^* \restriction [m^1_0, \omega)) \geq p$, and we have $p' \in \mathcal{J}$. Moreover $q \geq r^+ \geq r$ as required. So $\mathcal{J}$ is predense.

5. Part (α) follows immediately from the definition. For part (β) take into account that the m_i, $i < \omega$, in Definition 1.1(b)(γ) are unique, and hence $\langle m_i : i < \omega \rangle$ is already in the ground model where $\mathcal{C}^1$ exists.

Only parts (γ) and (δ) are not so short:

(γ) Suppose that p, q are incompatible in $\mathbb{A}(\mathcal{C}^1)$, and, in the first case, assume that $n^p \leq n^q$. The first subcase is $\bar{\mathfrak{c}}^p \restriction n^p \neq \bar{\mathfrak{c}}^q \restriction n^p$ or that they are equal, but $\bar{\mathfrak{c}}^q \restriction [n^p, n^q) \not\geq_{\rm full} \bar{\mathfrak{c}}^p \restriction [n^p, \infty)$. Then, of course, this is also a reason for incompatibility in $\mathbb{A}(\mathcal{C}^2)$. The second subcase is $\bar{\mathfrak{c}}^p \restriction n^p = \bar{\mathfrak{c}}^q \restriction n^p$ and $\bar{\mathfrak{c}}^q \restriction [n^p, n^q) \geq_{\rm full} \bar{\mathfrak{c}}^p \restriction [n^p, \infty)$, but for all $\bar{\mathfrak{c}} \in \mathcal{C}^1$, $\bar{\mathfrak{c}}^p \restriction [n^p, \omega) \not\leq_{\rm full} \bar{\mathfrak{c}} \restriction [n^p, \omega)$ or $\bar{\mathfrak{c}}^q \restriction [n^q, \omega) \not\leq_{\rm full} \bar{\mathfrak{c}} \restriction [n^q, \omega)$. By the premise $\bar{\mathfrak{c}}^p \leq_{\mathbf{q}} \bar{\mathfrak{c}}^q$, there $i(*)$ and $n_{i(*)}$ such that $\bar{\mathfrak{c}}^p \restriction [n_{i(*)}, \omega) \leq_{\rm full} \bar{\mathfrak{c}}^q \restriction [i(*), \omega)$, and $i(*), n_{i(*)} \geq n^p$. By forgetfulness, there are $m_1, m_2 > n^q$ and $\bar{\mathfrak{d}}$ such that $m_{\rm dn}(\mathfrak{c}^p_{m_1}) = m_{\rm dn}(\mathfrak{c}^q_{m_2})$ and

$$\bar{\mathfrak{d}} \in \Sigma(\bar{\mathfrak{c}}^p \restriction [n^q, m_1) \cap \Sigma(\bar{\mathfrak{c}}^q \restriction [n^q, m_2) \neq \emptyset.$$

Then $\bar{\mathfrak{c}} = (\bar{\mathfrak{c}}^q \restriction n^q{}^\frown \bar{\mathfrak{d}}{}^\frown \bar{\mathfrak{c}}^q \restriction [m_2, \omega) \in \mathcal{C}^1$ shows that p, q are compatible in $\mathbb{A}(\mathcal{C}^1)$. Contradiction. The case $n^q < n^p$ is similar.

(δ) Again, as in (γ), we may assume that $n^p \leq n^q$ and that we are in the second subcase. We show that if $(n^p, \bar{\mathfrak{c}}^p)$ and $(n^q, \bar{\mathfrak{c}}^q)$ are compatible in $\mathbb{A}(\mathcal{C}^2)$, then they are compatible in $\mathbb{A}(\mathcal{C}^1)$.

Since $\mathcal{C}^1$ is $\mathbf{q}$-directed, there is some $\bar{\mathfrak{c}} \in \mathcal{C}^1$ such that $\bar{\mathfrak{c}}^p, \bar{\mathfrak{c}}^q \leq \bar{\mathfrak{c}}$. By $\bar{\mathfrak{c}}^p \leq_{\mathbf{q}} \bar{\mathfrak{c}}$, there $i_p(*)$ and $n_{i_p(*)}$ such that $\bar{\mathfrak{c}}^p \restriction [n_{i_p(*)}, \omega) \leq_{\rm full} \bar{\mathfrak{c}} \restriction [i_p(*), \omega)$, and $i_p(*), n_{i_p(*)} \geq n^p$. By $\bar{\mathfrak{c}}^q \leq_{\mathbf{q}} \bar{\mathfrak{c}}$, there $i_q(*)$ and $n_{i_q(*)}$ such that $\bar{\mathfrak{c}}^p \restriction [n_{i_q(*)}, \omega) \leq_{\rm full} \bar{\mathfrak{c}} \restriction [i_q(*), \omega)$, and $n_{i_p(*)} \geq n^p$, w.l.o.g. $n_{i_q(*)} \geq n^q$.

By the premise there in some $(n, \bar{\mathfrak{d}}) \in \mathbb{A}(\mathcal{C}^2)$ such that $(n, \bar{\mathfrak{d}}) \geq (n^p, \bar{\mathfrak{c}}^p), (n^q, \bar{\mathfrak{c}}^q)$. So $\bar{\mathfrak{d}} \restriction [n^p, \infty) \geq_{\rm full} \bar{\mathfrak{c}}^p \restriction [n^p, \infty)$ and $\bar{\mathfrak{d}} \restriction [n^q, \infty) \geq_{\rm full} \bar{\mathfrak{c}}^q \restriction [n^q, \infty)$. So in particular there is some $m < \omega$ such that $\bar{\mathfrak{d}} \restriction [n^p, m) \geq_{\rm full} \bar{\mathfrak{c}}^p \restriction [n^p, n_{i_q(*)})$ and $\bar{\mathfrak{d}} \restriction [n^q, m) \geq_{\rm full} \bar{\mathfrak{c}}^q \restriction [n^q, n_{i_q(*)})$. Then $((\bar{\mathfrak{d}} \restriction m){}^\frown \bar{\mathfrak{c}}) \in \mathcal{C}^1$ since $\mathcal{C}^1$ is closed under $\equiv_{\mathbf{q}}$

in $\mathbf{V}$ and all finite sections of $\bar{\mathfrak{d}}$ are in $\mathbf{V}$. $(m, (\bar{\mathfrak{d}} \restriction m)\hat{}\bar{\mathfrak{c}} \restriction [i_q(*), \omega))$ is a stronger in $\mathbb{A}(\mathcal{C}^1)$ than $(n^p, \bar{\mathfrak{c}}^p)$ and $(n^q, \bar{\mathfrak{c}}^q)$.

6. For $\mathcal{C}^1 \subseteq \mathcal{C}^2$ being $\mathbf{q}$-pre-directed, we can choose $D_1 \subseteq D_2$. In the limit we take the union of the proper filters, which is again a proper filter, i.e., all intersections of finitely many of its members are infinite and members of the filter. Any finitely many elements from $\bigcup_{\alpha<\lambda} \mathcal{C}^\alpha$ have an upper bound whose support is in the filter.

For the $\mathcal{C}^\alpha$'s being $\mathbf{q}$-directed and increasing, it is clear that their union is $\mathbf{q}$-directed. $\square$

Definition 1.15. *Let $\mathbf{q}$ be a forgetful pcrf.*

1. *Let $\mathcal{C}$ be $\mathbf{q}$-directed, $\bar{\mathfrak{d}} \in \mathcal{C}$ and $\mathcal{J} \subseteq \mathbb{A}_{\mathcal{C}}(\bar{\mathfrak{d}})$ be a maximal antichain of $\mathbb{A}(\mathcal{C})$. By induction on the ordinal α we define when $\mathrm{rk}_{\mathbf{q}}(n, m, \bar{\mathfrak{c}}, \bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C}) = \alpha$ for $\bar{\mathfrak{c}} \in \mathbf{Q}_{\mathcal{C}}(\bar{\mathfrak{d}})$ and $n \le m < \omega$.*

   Case: $\alpha = 0$. $\mathrm{rk}_{\mathbf{q}}(n, m, \bar{\mathfrak{c}}, \bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C}) = 0$ iff $(n, \bar{\mathfrak{c}}) \in \mathbb{A}(\mathcal{C})$ is above some member of $\mathcal{J}$.

   Case: $\alpha > 0$. $\mathrm{rk}_{\mathbf{q}}(n, m, \bar{\mathfrak{c}}, \bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C}) = \alpha$ iff

   (a) *for no $\beta < \alpha$ we have $\mathrm{rk}_{\mathbf{q}}(n, m, \bar{\mathfrak{c}}, \bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C}) = \beta$,*

   (b) *for all $p' = (m', \bar{\mathfrak{d}}') \in \mathcal{J}$, $\beta < \alpha$, exists $\bar{\mathfrak{c}}'$ such that*

      (α) *$\bar{\mathfrak{c}}' \in \mathbf{Q}_{\mathcal{C}}(\bar{\mathfrak{d}})$ (hence $\bar{\mathfrak{c}}' \le_{\mathbf{q}} \bar{\mathfrak{d}}'$ and vice versa and same with $\bar{\mathfrak{c}}$, $\bar{\mathfrak{d}}$, because all of them are $\equiv_{\mathbf{q}} \bar{\mathfrak{d}}$),*

      (β) *$\bar{\mathfrak{c}} \restriction m = \bar{\mathfrak{c}}' \restriction m$,*

      (γ) *$(n, \bar{\mathfrak{c}}')$ and $(m', \bar{\mathfrak{d}}')$ are incompatible in $\mathbb{A}_{\mathcal{C}}(\bar{\mathfrak{d}})$, equivalently (—by (b)($\alpha$) and Claim 1.14(5) ($\delta$)—) in $\mathbb{A}(\mathcal{C})$,*

      (δ) *$\beta \le \mathrm{rk}_{\mathbf{q}}(n, m+1, \bar{\mathfrak{c}}', \bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C})$.*

2. $\mathrm{rk}_{\mathbf{q}}(\bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C}) = \bigcup\{\mathrm{rk}_{\mathbf{q}}(n, m, \bar{\mathfrak{c}}, \bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C}) + 1 \,:\, \bar{\mathfrak{c}} \in \mathbf{Q}_{\mathcal{C}}(\bar{\mathfrak{d}}), n \le m < \omega\}$.

3. *We say $\mathcal{C}$ is $\mathbf{q}$-nice if for all $\bar{\mathfrak{d}} \in K$, $\mathcal{J} \subseteq \mathbb{A}_{\mathcal{C}}(\bar{\mathfrak{d}})$ that are predense in $\mathbb{A}(\mathcal{C})$,* $\mathrm{rk}_{\mathbf{q}}(\bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C}) < \omega_1$. *Remark: Since $\mathbb{A}(\mathcal{C})$ is σ-centred and since countable sets have $\le_{\mathbf{q}}$ upper bounds, there are such $\bar{\mathfrak{d}}$'s and $\mathcal{J}$'s.*

   $\forall \mathcal{J}\, \mathrm{rk}_{\mathbf{q}}(\bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C}) < \omega_1$ is absolute: $(\forall \mathcal{J} \subseteq \mathbb{A}_{\mathcal{C}}(\bar{\mathfrak{d}}))(\exists \alpha < \omega_1)(\mathrm{rk}_{\mathbf{q}}(\bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C}) = \alpha)$ is Π^1_2 and hence Shoenfield's absoluteness theorem applies. Also quantification over all $\bar{\mathfrak{d}}$ is possible. But if $\mathcal{C}$ is expanded, the rank can become infinite.

4. *We say* $\mathcal{J} \subseteq \mathbb{A}_{\mathcal{C}}(\bar{\mathfrak{d}})$ *is an explicitly predense set (or maximal antichain in* $\mathbb{A}(\mathcal{C})$*) if its members are pairwise incompatible and if* $\mathrm{rk}_{\mathbf{q}}(\bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C}) < \omega_1$.

Now, with the help of the rank function, we establish criteria for $\mathcal{J}$ to stay predense in $\mathbb{A}(\mathcal{C})$ in any extension of the universe and under any extension of $\mathcal{C}$. As usual, we write $\mathbf{V}^{\mathbb{P}}$ for $\mathbf{V}[G]$ with an arbitrary $\mathbb{P}$-generic G over $\mathbf{V}$.

Claim 1.16. *Assume*

(a) $\mathcal{C}$ *is a* $\mathbf{q}$*-directed family.*

(b) $\bar{\mathfrak{d}} \in \mathcal{C}$, *and* $\mathcal{J} \subseteq \mathbb{A}_{\mathcal{C}}(\bar{\mathfrak{d}})$ *is predense in* $\mathbb{A}(\mathcal{C})$.

Then the following six conditions are equivalent:

(α) *In* $\mathbf{V}^{\mathrm{Cohen}}$ *there is some* $\bar{\mathfrak{d}}'$ *such that* $\bar{\mathfrak{d}} \leq_{\mathbf{q}} \bar{\mathfrak{d}}' \in \mathbf{Q_q}^{\mathbf{V}^{\mathrm{Cohen}}}$, $\mathcal{C} \cup \{\bar{\mathfrak{d}}'\}$ *is a* $\mathbf{q}$*-pre-directed family, and for some* $n < \omega$, $(n, \bar{\mathfrak{d}}')$ *is incompatible in* $\mathbb{Q}(\mathcal{C} \cup \{\bar{\mathfrak{d}}'\})$ *with every member of* $\mathcal{J}$.

(β) *Like (α) for some atomless forcing notion* $\mathbb{P}$.

(β′) *For some atomless forcing notion* $\mathbb{P}$, *in* $\mathbf{V}^{\mathbb{P}}$ *there are some* $\bar{\mathfrak{d}}'$ *and some* n *and some* $\mathbf{q}$*-directed* $\mathcal{C}'$ *such that* $\mathbf{Q}_{\mathcal{C}}(\bar{\mathfrak{d}}) \subseteq \mathcal{C}'$, *such that* $\bar{\mathfrak{d}} \leq_{\mathbf{q}} \bar{\mathfrak{d}}' \in \mathcal{C}'$, *and in* $\mathbb{A}(\mathcal{C}')$ *the condition* $(n, \bar{\mathfrak{d}}')$ *is incompatible with every member of* $\mathcal{J}$.

(γ) *For some atomless forcing notion* $\mathbb{P}$, *in* $\mathbf{V}^{\mathbb{P}}$ *there is a strongly directed* $\mathcal{C}' \supseteq \mathcal{C}$, *such that in* $\mathbb{A}(\mathcal{C}')$, *the set* $\mathcal{J}$ *is not a maximal antichain.*

(δ) $\mathrm{rk}_{\mathbf{q}}(\bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C}) = \infty$.

(ε) $\mathrm{rk}_{\mathbf{q}}(\bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C}) \geq \omega_1$.

Proof. The proof is via a circle $(\varepsilon) \Rightarrow (\alpha) \Rightarrow (\beta) \Rightarrow (\beta') \Rightarrow (\gamma) \Rightarrow (\delta) \Rightarrow (\varepsilon)$.

$(\varepsilon) \Rightarrow (\alpha)$. $\mathrm{rk}_{\mathbf{q}}(\bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C}) \geq \omega_1$. We choose $\bar{\mathfrak{d}}'' \restriction m$ by induction on m. Let $\mathcal{J} = \{p'_k : k \in \omega\}$ be an enumeration of $\mathcal{J}$. We take $n, m, \bar{\mathfrak{c}}$ such that $\mathrm{rk}_{\mathbf{q}}(n, m, \bar{\mathfrak{c}}, \bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C}) \geq \omega_1$. Since $\omega \times \omega \times \mathbf{Q}_{\mathcal{C}}(\bar{\mathfrak{d}})$ is countable and since ω_1 is regular, there is such a triple. Let $\mathbb{P}$ be the forcing adding a Cohen real. Suppose that $m_0, \bar{\mathfrak{c}}^0, p'_k$ are given and that $\omega_1 = \mathrm{rk}_{\mathbf{q}}(n, m_0, \bar{\mathfrak{c}}^0, \bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C})$. Let $D_{p'_k, m_0, \bar{\mathfrak{c}}^0}$ be the set of $\bar{\mathfrak{c}}^1$ such that

(1) $\bar{\mathfrak{c}}^1 \in \mathbf{Q}_{\mathcal{C}}(\bar{\mathfrak{d}})$,

(2) $\bar{\mathfrak{c}}^1 \restriction m_0 = \bar{\mathfrak{c}}^0 \restriction m_0$,

(3) $(n, \bar{\mathfrak{c}}^1)$ and p'_k are incompatible in $\mathbb{A}_{\mathcal{C}}(\bar{\mathfrak{d}})$ and the incompatibility is witnessed below $m_0 + \ell$ for some ℓ and (— so, despite forgetfulness, there are

also many elements outside $\Sigma(\bar{\mathfrak{c}}^1 \restriction [k, m))$ that are in $\mathbb{A}_{\mathcal{C}}(\bar{\mathfrak{d}})$ and incompatible with p'_k—)

(4) $\forall \beta < \omega_1$, $\beta \leq \mathrm{rk}_{\mathbf{q}}(n, m_0 + \ell, \bar{\mathfrak{c}}^1, \bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C})$.

Reading definition 1.15(b) ℓ times shows that $D_{p'_k, m_0, \bar{\mathfrak{c}}^0}$ is is dense in the Cohen forcing. So, if G is $\mathbb{A}_{\mathcal{C}}(\bar{\mathfrak{d}})$-generic over $\mathbf{V}$ and $\bar{\mathfrak{d}}' = \bigcup\{\bar{\mathfrak{c}} \restriction n : (n, \bar{\mathfrak{c}}) \in G\}$, then $(n, \bar{\mathfrak{d}}')$ is incompatible in $\mathbb{A}(\mathcal{C} \cup \{\bar{\mathfrak{d}}'\})$ with every member of $\mathcal{J}$.

$(\beta) \Rightarrow (\beta')$: by Claim 1.14 Part 5 (γ): Incompatibility of $\bar{\mathfrak{d}}'$ with every member of $\mathcal{J}$ is the same in $\mathbb{A}(\mathcal{C} \cup \{\bar{\mathfrak{d}}'\})$ and in $\mathbb{A}(\mathcal{C}')$, because every member of $\mathcal{J}$ is $\leq_{\mathbf{q}} \bar{\mathfrak{d}}'$.

$(\beta') \Rightarrow (\gamma)$: We take a model from (β') and extend $\mathcal{C}'$ from that model to a strongly directed $\mathcal{C}''$. This is possible because $\leq_{\mathbf{q}}$ increasing countable chains have an upper bound in $\mathbf{Q}$. Since $\mathcal{C}'$ is directed and all intermediate extensions can chosen to be directed, it is sufficient to add limits to chains instead of upper bounds to countable subsets.

$(\gamma) \Rightarrow (\delta)$ $\mathrm{rk}_{\mathbf{q}}(n, m, \bar{\mathfrak{d}}' \restriction m, \bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C})$ is witness to a descending chain in the ranks. Since $K \subseteq \mathcal{H}(\aleph_0)$ also for all new $\bar{\mathfrak{d}}'$'s, every $\bar{\mathfrak{d}}' \restriction m \in \mathbf{V}$. □

Conclusion 1.17. *1. If $\mathbb{P}$ is the forcing for adding $2^{\aleph_0}$ Cohen reals, then for any $\mathbf{q}$-directed $\mathcal{C}^1 \in \mathbf{V}$, in $\mathbf{V}^{\mathbb{P}}$ we can extend $\mathcal{C}^1$ to a $\mathbf{q}$-nice $\mathcal{C}^2$.*

2. Assume $\mathbb{P}_1 \lessdot \mathbb{P}_2$ and $\Vdash_{\mathbb{P}_1}$ "$\underset{\sim}{\mathcal{C}}^1$ is $\mathbf{q}$-nice" and $\Vdash_{\mathbb{P}_2}$ "$\underset{\sim}{\mathcal{C}}^2$ is $\mathbf{q}$-directed and extends $\underset{\sim}{\mathcal{C}}^1$". Then $\mathbb{P}_1 * \mathbb{A}(\underset{\sim}{\mathcal{C}}^1) \lessdot \mathbb{P}_2 * \mathbb{A}(\underset{\sim}{\mathcal{C}}^2)$.

3. If $\mathbb{P}_2/\mathbb{P}_1$ contains as a complete subforcing the forcing for adding $2^{\aleph_0}$ Cohen reals, then there is $\underset{\sim}{\mathcal{C}}^3$ such that $\Vdash_{\mathbb{P}_2}$ "$\mathcal{C}^3$ is $\mathbf{q}$-nice and extends $\underset{\sim}{\mathcal{C}}^1$", and $\mathbb{P}_1 * \mathbb{A}(\underset{\sim}{\mathcal{C}}^1) \lessdot \mathbb{P}_2 * \mathbb{A}(\underset{\sim}{\mathcal{C}}^3)$.

Proof. 1. Let $\langle (\bar{\mathfrak{d}}_\alpha, \mathcal{J}_\alpha) : \alpha < 2^{\aleph_0} \rangle$ be an enumeration such that for all $\beta < 2^{\aleph_0}$, $\langle (\bar{\mathfrak{d}}_\alpha, \mathcal{J}_\alpha) : \alpha \geq \beta \rangle$ enumerates among others all of $\bigcup\{K^{\mathbf{V}[\bar{\mathfrak{e}}]} \times \mathcal{P}(\mathbb{A}_{\mathcal{C}) \cup \mathrm{range}\, \bar{\mathfrak{e}}}(\bar{\mathfrak{d}}) : \bar{\mathfrak{e}} \in V[\mathrm{Cohen}_\beta] \cap (\omega^\omega)^\omega, \bar{\mathfrak{d}} \in \mathcal{C} \cup \mathrm{range}\, \bar{\mathfrak{e}}\}$. Then for step $\alpha < 2^{\aleph_0}$ of an iterative construction take $(\bar{\mathfrak{d}}_\alpha, \mathcal{J}_\alpha)$ check whether (α) of claim 1.16 is true for $(\bar{\mathfrak{d}}_\alpha, \mathcal{J}_\alpha, \mathcal{C}_\alpha)$, and if so, then add by Cohen forcing $\bar{\mathfrak{d}}'$ as there, and define $\mathcal{C}_{\alpha+1} = \mathcal{C}_\alpha \cup \{\bar{\mathfrak{d}}'\}$.

2. Similar to [10, Claim 4.7] or to [11, Section 5, Claim 5.6]. For completeness, we give the proof: Since we may think of $\mathbb{P}_2$ as $\mathbb{P}_2/\mathbb{P}_1$, we can replace $\mathbb{P}_1$ by the trivial forcing. Then $\mathcal{C}^1$ is in the ground model. Let G be a $\mathbb{P}_2$-generic filter and let $\mathcal{C}^2 = \underset{\sim}{\mathcal{C}}^2[G]$. If $p, q \in \mathbb{A}(\mathcal{C}^1)$ and $p \leq q$, then they are also in $\mathbb{A}(\mathcal{C}^2)$ in this order. The same holds for incompatibility by Claim 1.14(5). Now we need to show that maximal antichains in $\mathbb{A}(\mathcal{C}^1)$ stay maximal in $\mathbb{A}(\mathcal{C}^2)$: For this take a

maximal antichain $\mathcal{J}$. It is countable and a subset of $\mathbb{A}_{\mathcal{C}^1}(\bar{\mathfrak{d}})$ for some suitable $\bar{\mathfrak{d}} \in \mathbf{V}$. Now we have by $\mathcal{C}^1$ being nice that $\mathrm{rk}(\bar{\mathfrak{d}}, \mathcal{J}, \mathcal{C}^1) < \omega_1$. This is absolute and holds in $V[G]$ as well. So no member of $\mathbb{A}(\mathcal{C}^2)$ can be orthogonal to a member of $\mathcal{J}$.

3. This is just the combination of the first two parts. □

Definition 1.18. *Assume that $\mathcal{C}$ is $\mathbf{q}$-directed. Let $\underset{\sim}{G}_{\mathbb{A}(\mathcal{C})}$ be a name for the $\mathbb{A}(\mathcal{C})$-generic filter G. $\underset{\sim}{\bar{\mathfrak{c}}}_{\mathbb{A}(\mathcal{C})}$ is the $\mathbb{A}(\mathcal{C})$-name*

$$\bigcup\{\bar{\mathfrak{c}} \restriction n \,:\, (n, \bar{\mathfrak{c}}) \in \underset{\sim}{G}_{\mathbb{A}(\mathcal{C})}\}.$$

Claim 1.19. *Assume that $\mathcal{C}$ is $\mathbf{q}$-directed.*

1. $\Vdash_{\mathbb{A}(\mathcal{C})}$ "$\underset{\sim}{\bar{\mathfrak{c}}} \in \mathbf{Q}$ is a $\leq_{\mathbf{q}}$ -upper bound of $\mathcal{C}$".

2. If $\mathcal{C}$ is a $\mathbf{q}$-nice family, then $\underset{\sim}{\bar{\mathfrak{c}}}$ is a generic real for $\mathbb{A}(\mathcal{C})$, that is, for $G \subseteq \mathbb{A}(\mathcal{C})$ generic over $\mathbf{V}$, letting $\bar{\mathfrak{c}}^* = \underset{\sim}{\bar{\mathfrak{c}}}[G]$, we have

$$G = \{p \in \mathbb{Q}(\mathcal{C}) \,:\, \bar{\mathfrak{c}}^p \restriction n^p = \bar{\mathfrak{c}}^* \restriction n^p \text{ and } \bar{\mathfrak{c}}^p \restriction [n^p, \infty) \leq_{\text{full}} \bar{\mathfrak{c}}^* \restriction [n^p, \omega)\}. \tag{1.6}$$

Proof. 1. For all $\bar{\mathfrak{d}} \in \mathcal{C}$, the set $\{(n^p, \bar{\mathfrak{c}}^p) \in \mathbb{A}(\mathcal{C}) \,:\, \bar{\mathfrak{c}}^p \geq_{\mathbf{q}} \bar{\mathfrak{d}}\}$ is open and dense in $\mathbb{A}(\mathcal{C})$, because $\mathcal{C}$ is $\mathbf{q}$-directed, and because $\bar{\mathfrak{c}}^p \geq_{\mathbf{q}} \bar{\mathfrak{d}}$ and $(n^q, \bar{\mathfrak{c}}^q) \geq (n^p, \bar{\mathfrak{c}}^p)$ together implies $\bar{\mathfrak{c}}^q \geq_{\mathbf{q}} \bar{\mathfrak{d}}$.

2. First we show that G is a subset of the right hand side. Let $p \in G$. Then $\bar{\mathfrak{c}}^p \restriction n^p \lhd \bar{\mathfrak{c}}^*$. Also for every $m \geq n^p$ there is some $q \in G$ such that $n^q \geq m$ and $q \not\perp p$ hence $\exists m' \geq m$ $\bar{\mathfrak{c}}^p \restriction [n^p, m') \leq_{\text{full}} \bar{\mathfrak{c}}^q \restriction [n^p, n^q) = \bar{\mathfrak{c}}^* \restriction [n^p, n^q)$. Hence $\bar{\mathfrak{c}}^p \restriction [n^p, \omega) \leq_{\text{full}} \bar{\mathfrak{c}}^* \restriction [n^p, \omega)$.

Now for the other direction. Any two elements of the right hand side have $\bar{\mathfrak{c}}^*$ as a common stronger element. Hence the right hand side is directed and hence G cannot be a proper subset of it. □

Claim 1.20. *Assume*

(a) $\mathbf{q}$ is a pcrf,

(b) $\mathbb{P}_0 \lessdot \mathbb{P}_2$,

(c) for $\ell = 0, 2$ we have that $\underset{\sim}{\mathcal{C}}^\ell$ is a $\mathbb{P}_\ell$-name of a $\mathbf{q}$-nice family,

(d) for $\ell = 0, 2$ let $\mathbb{P}_{\ell+1} = \mathbb{P}_\ell * \mathbb{A}(\underset{\sim}{\mathcal{C}}^\ell)$, and let $\underset{\sim}{\bar{\mathfrak{c}}}^\ell$ be a $\mathbb{P}_{\ell+1}$-name of the generic for $\mathbb{A}(\underset{\sim}{\mathcal{C}}^\ell)$,

(e) $\Vdash_{\mathbb{P}_2}$ "$\underaccent{\tilde}{\mathcal{C}}^0 \subseteq \underaccent{\tilde}{\mathcal{C}}^2$".

Then

(α) $\mathbb{P}_2 \Vdash \underaccent{\tilde}{\bar{\mathfrak{c}}}^2$ *is a generic for* $\mathbb{A}(\underaccent{\tilde}{\mathcal{C}}^0)$.

(β) $\mathbb{P}_1 \lessdot \mathbb{P}_3$.

Proof. See 1.17 (2). $\mathbb{P}_0 * \mathbb{A}(\underaccent{\tilde}{\mathcal{C}}^0) \lessdot \mathbb{P}_2 * \mathbb{A}(\underaccent{\tilde}{\mathcal{C}}^0) \lessdot \mathbb{P}_2 * \mathbb{A}(\underaccent{\tilde}{\mathcal{C}}^2)$. □

2. Forcing bigness notions

See "Vive la différence III" [12] for a connection to model theory.

Definition 2.1. *1. A forcing bigness notion is a class* Γ*, partially ordered by* $\leq_\Gamma$*, such that for all* $(\mathbb{P}_0, \mathbb{P}_0^+, \underaccent{\tilde}{\eta}) \in \Gamma$ *we have*

(a) $\mathbb{P}_0 \lessdot \mathbb{P}_0^+$ *are forcing notions,*

(b) $\underaccent{\tilde}{\eta}$ *is a* $\mathbb{P}_0^+$*-name of a sequence of ordinals (usually* $\in {}^\omega\omega$*),*

(c) *if* $\mathbb{P} \lessdot \mathbb{P}_0$ *and* $\mathbb{P}_0^+ \lessdot \mathbb{P}^+$*, then* $(\mathbb{P}, \mathbb{P}^+, \underaccent{\tilde}{\eta}) \in \Gamma$.

If $(\mathbb{P}_0, \mathbb{P}_0^+, \underaccent{\tilde}{\eta}) \in \Gamma$*, we may say that* $(\mathbb{P}_0, \mathbb{P}_0^+, \underaccent{\tilde}{\eta})$ *is* Γ*-big.*

2. $\leq_\Gamma$ *is a quasi order and* $\lessdot_\Gamma \supseteq \leq_\Gamma$ *where* $\lessdot_\Gamma$ *is the following (quasi) partial order:* $(\mathbb{P}_1, \mathbb{P}_1^+, \underaccent{\tilde}{\eta}_1) \lessdot_\Gamma (\mathbb{P}_2, \mathbb{P}_2^+, \underaccent{\tilde}{\eta}_2)$ *means that both triples are in* Γ*, and* $\mathbb{P}_1 \lessdot \mathbb{P}_2$ *and* $\mathbb{P}_1^+ \lessdot \mathbb{P}_2^+$ *and* $\underaccent{\tilde}{\eta}_1 = \underaccent{\tilde}{\eta}_2$ *and such that the following diagramme is exact, which means, that for any* $p_1^+ \in \mathbb{P}_1^+$*,* $p_2 \in \mathbb{P}_2$*,*

$$\sup_{\mathbb{P}_1}\{p_1 \in \mathbb{P}_1 : p_1 \perp_{\mathbb{P}_1^+} p_1^+\} = \sup_{\mathbb{P}_2}\{p_2 \in \mathbb{P}_2 : p_2 \perp_{\mathbb{P}_2^+} p_1^+\}. \tag{2.1}$$

$$\begin{tikzcd}
\mathbb{P}_1^+ \arrow[r, "\lessdot"] & \mathbb{P}_2^+ \\
\mathbb{P}_1 \arrow[u, "\lessdot"] \arrow[r, "\lessdot"] & \mathbb{P}_2 \arrow[u, "\lessdot"]
\end{tikzcd}$$

3. We say Γ *is a c.c.c. forcing bigness notion, if in addition to the above mentioned properties the first and second components of all triples in* Γ *are c.c.c forcing notions and if*

(d) $\langle(\mathbb{P}_i, \mathbb{P}_i^+, \underaccent{\tilde}{\eta}) : i < \delta\rangle \in \Gamma$ *is continuously increasing in* $\leq_\Gamma$*, then* $(\bigcup\{\mathbb{P}_i : i < \delta\}, \bigcup\{\mathbb{P}_i^+ : i < \delta\}, \underaccent{\tilde}{\eta}) \in \Gamma$ *and is* $\leq_\Gamma$*-above all* $(\mathbb{P}_i, \mathbb{P}_i^+, \underaccent{\tilde}{\eta})$.

4. For a c.c.c. forcing bigness notion Γ *we say that* Γ *has amalgamation iff: If* $(\mathbb{P}_1, \mathbb{P}_1^+, \underaccent{\tilde}{\eta}) \in \Gamma$ *and* $\mathbb{P}_1 \lessdot \mathbb{P}_2$ *and* $\mathbb{P}_2/\mathbb{P}_1$ *satisfies the Knaster property, then we can find* $\mathbb{P}_2^+$ *[satisfying the c.c.c.] and* f *such that* *(α)* $\mathbb{P}_1^+ \lessdot \mathbb{P}_2^+$,

(β) *f is a complete embedding of $\mathbb{P}_2$ into $\mathbb{P}_2^+$ over $\mathbb{P}_1$,*

(γ) $(f(\mathbb{P}_2), \mathbb{P}_2^+, \underset{\sim}{\eta}) \in \Gamma$,

(δ) $(\mathbb{P}_1, \mathbb{P}_1^+, \underset{\sim}{\eta}) \leq_\Gamma (f(\mathbb{P}_2), \mathbb{P}_2^+, \underset{\sim}{\eta})$.

5. For a c.c.c. forcing bigness notion Γ we say that Γ has free amalgamation iff: If $(\mathbb{P}_1, \mathbb{P}_1^+, \underset{\sim}{\eta}) \in \Gamma$ and $\mathbb{P}_1 \lessdot \mathbb{P}_2$ and $\mathbb{P}_2/\mathbb{P}_1$ satisfies the Knaster property, then we can choose in 4. $\mathbb{P}_2^+ = \mathbb{P}_1^+ \times_{\mathbb{P}_1} \mathbb{P}_2$ [satisfying the c.c.c.] such that

(α) $\mathbb{P}_1^+ \lessdot \mathbb{P}_2^+$,

(β) *id is a complete embedding of $\mathbb{P}_2$ into $\mathbb{P}_2^+$ over $\mathbb{P}_1$,*

(γ) $(\mathbb{P}_2, \mathbb{P}_2^+, \underset{\sim}{\eta}) \in \Gamma$.

For a definition of $\mathbb{P}_1^+ \times_{\mathbb{P}_1} \mathbb{P}_2$ see [8].

6. We say Γ has the $(< \lambda)$-LS (Löwenheim Skolem) property when the following holds: If $(\mathbb{P}, \mathbb{P}^+, \underset{\sim}{\eta}) \in \Gamma$ and $Y \subseteq \mathbb{P}^+$, $|Y| < \lambda$ (or alternatively, for stationarily many $Y \in [\mathbb{P}^+]^{<\lambda}$) we have: There is $(\mathbb{P}_1, \mathbb{P}_1^+, \underset{\sim}{\eta})$ such that,

(α) $(\mathbb{P}_1, \mathbb{P}_1^+, \underset{\sim}{\eta}) \in \Gamma$,

(β) $|\mathbb{P}_1^+| < \lambda$,

(γ) $\mathbb{P} \cap Y \subseteq \mathbb{P}_1 \lessdot \mathbb{P}$ *and*

(δ) $Y \subseteq \mathbb{P}_1^+ \lessdot \mathbb{P}^+$.

7. Γ is a Knaster forcing bigness notion iff it is a c.c.c. forcing bigness notion and in addition, if $(\mathbb{P}, \mathbb{P}^+, \underset{\sim}{\eta}) \in \Gamma$ then $\mathbb{P}$ and $\mathbb{P}^+$ satisfy the Knaster condition.

Observation 2.2. *$\lessdot_\Gamma$ is a partial order.*

Definition 2.3. *(1) We define the following Γ:*

- *(a)* $\Gamma_{\text{new}} = \{(\mathbb{P}, \mathbb{P}^+, \underset{\sim}{\eta}) : \mathbb{P} \lessdot \mathbb{P}^+$ *are c.c.c. forcing notions and such that* $\Vdash_{\mathbb{P}^+} \underset{\sim}{\eta} \in ({}^\omega\omega)^{\mathbf{V}[G_{\mathbb{P}^+}]} \setminus \mathbf{V}[G_\mathbb{P}]\}$.
- *(b)* $\Gamma_{\text{Cohen}} = \{(\mathbb{P}, \mathbb{P}^+, \underset{\sim}{\eta}) \in \Gamma_{\text{new}} : \underset{\sim}{\eta}$ *is forced by $\mathbb{P}^+$ to be a Cohen real over* $\mathbf{V}^\mathbb{P}\}$.
- *(c)* $\Gamma_{\text{ud}} = \{(\mathbb{P}, \mathbb{P}^+, \underset{\sim}{\eta}) \in \Gamma_{\text{new}} : \underset{\sim}{\eta}$ *is forced by $\mathbb{P}^+$ not to be dominated by any $\nu \in \mathbf{V}^{[G_\mathbb{P}]}\}$. η is dominted by ν, abbreviated $\eta \leq^* \nu$, if* $(\exists n)(\forall k \geq n)(\eta(k) \leq \nu(k))$.

(d) $\Gamma_{\rm tow} = \{(\mathbb{P}, \mathbb{P}^+, \underset{\sim}{\eta}) \in \Gamma_{\rm new} : \mathbb{P} \lessdot \mathbb{P}^+$ *and* $\underset{\sim}{\eta} \in ({}^\omega 2)^{\mathbf{V}[G_{\mathbb{P}^+}]}$ *are such that* $\eta^{-1}\{1\}$ *is infinite and is forced by* $\mathbb{P}^+$ *not to contain any infinite subset of* ω *which belongs to* $\mathbf{V}^{\mathbb{P}}\}$

(2) *We define when a name* $\underset{\sim}{\eta}$ *is* Γ*-big.*

(a) *We say* $\underset{\sim}{\eta}$ *is* $\Gamma_{\rm new}$*-big over* $\nu \in \mathbf{V}[\mathbb{P}]$ *if* $\Vdash_{\mathbb{P}^+} \underset{\sim}{\eta} \notin \mathbf{V}[\nu]$.

(b) *We say* $\underset{\sim}{\eta}$ *is* $\Gamma_{\rm Cohen}$*-big over* $\nu \in \mathbf{V}[\mathbb{P}]$ *if* $\Vdash_{\mathbb{P}^+} \underset{\sim}{\eta}$ *Cohen-generic over* $\mathbf{V}[\nu]$.

(c) *We say* $\underset{\sim}{\eta}$ *is* $\Gamma_{\rm ud}$*-big over* $\nu \in \mathbf{V}[\mathbb{P}]$ *if* $\Vdash_{\mathbb{P}^+} \underset{\sim}{\eta} \not\leq^* \nu$.

(d) *We say* $\underset{\sim}{\eta}$ *is* $\Gamma_{\rm tow}$*-big over* $\nu \in \mathbf{V}[\mathbb{P}]$ *if* $\Vdash_{\mathbb{P}^+} \underset{\sim}{\eta}^{-1}[\{1\}]$ *does not contain an infinite subset from* $\mathbf{V}[\nu]$.

Definition 2.4. *(1) For any pseudo creature forcing* $\mathbf{q}$ *we write*

$$\Gamma_{\mathbf{q}} = \{(\mathbb{P}, \mathbb{P}^+, \underset{\sim}{\eta}) \in \Gamma_{\rm new} : \underset{\sim}{\eta} \in \mathbf{Q}^{\mathbf{V}[\mathbb{P}^+]} \wedge \Vdash_{\mathbb{P}^+} \text{“for no } \bar{\mathfrak{c}} \in \mathbf{Q}^{\mathbf{V}[\mathbb{P}]} \text{ we have } \underset{\sim}{\eta} \leq_{\mathbf{q}} \bar{\mathfrak{c}}\text{”}\}. \tag{2.2}$$

We can allow $\mathbf{q}$ *to be a* $\mathbb{P}$*-name.*

(2) *We say* $\underset{\sim}{\eta}$ *is* $\Gamma_{\mathbf{q}}$*-big over* $\nu \in \mathbf{V}[\mathbb{P}]$ *if for no* $\bar{\mathfrak{c}} \in \mathbf{V}[\nu]$ $\Vdash_{\mathbb{P}^+} \underset{\sim}{\eta} \leq_{\mathbf{q}} \bar{\mathfrak{c}}$.

Claim 2.5. *The* Γ*'s from Definitions 2.3 and 2.4 together with the partial ordering* $\lessdot_\Gamma$ *are c.c.c. forcing bigness notions with amalgamation. Moreover, if* $(\forall \mu < \lambda)(\mu^{\aleph_0} < \lambda)$ *then they have the* $(< \lambda)$*-LS.*

Proof. (a) We need to check all the items in Definition 2.1 parts 1. and 3. and 4. 1.(a),(b),(c) follow directly from the definition of $\Gamma_{\rm new}$.

Now for part 3: we first need to show that the union of c.c.c. forcings $\mathbb{P}_i$, $i < \delta$, is again c.c.c. Since the chain is $\lessdot$ increasing, we can fix complete projections for $i < j < \delta$ $\mathrm{pr}_{j,i} : \mathbb{P}_j \to \mathbb{P}_i$ (see [1] for the characterization of $\lessdot$ via complete projections) that form a commutative system. Let $\{p_\alpha : \alpha < \omega_1\} \subseteq \bigcup\{\mathbb{P}_i : i < \delta\}$. We show that this is not an antichain. Suppose that $p_\alpha \in \mathbb{P}_{i(\alpha)}$ and such that for $\alpha < \beta$, $i(\alpha) \leq i(\beta)$. Since $\mathbb{P}_0$ has the c.c.c, there are $\alpha < \beta$ such that $\mathrm{pr}_{i(\alpha),0}(p_\alpha) ||_0 \, \mathrm{pr}_{i(\beta),0}(p_\beta)$. But then $p_\alpha ||_\beta \, p_\beta$ and hence $p_\alpha ||_\delta \, p_\beta$.

Next we have to show that $\mathbb{P}_i \lessdot \bigcup\{\mathbb{P}_i : i < \delta\}$. This is again easy chasing arrows.

Now we need to show that $\bigcup\{\mathbb{P}_i : i<\delta\} \lessdot \bigcup\{\mathbb{P}_i^+ : i<\delta\}$ and that

$$\begin{tikzcd}
\mathbb{P}_1^+ \arrow[r, "\lessdot"] & \mathbb{P}_2^+ \arrow[r, "\lessdot"] & \cdots \arrow[r, "\lessdot"] & \bigcup\{\mathbb{P}_i^+ : i<\delta\} \\
\mathbb{P}_1 \arrow[u, "\lessdot"] \arrow[r, "\lessdot"] & \mathbb{P}_2 \arrow[u, "\lessdot"] \arrow[r, "\lessdot"] & \cdots \arrow[r, "\lessdot"] & \bigcup\{\mathbb{P}_i : i<\delta\} \arrow[u, "\lessdot"]
\end{tikzcd} \tag{2.3}$$

together with its complete embeddings and their inverse projections $\mathrm{pr}_{j,i}\colon \mathbb{P}_j \to \mathbb{P}_i$, $\mathrm{pr}_{j^+,i^+}\colon \mathbb{P}_j^+ \to \mathbb{P}_i^+$, $\mathrm{pr}_{i^+,i}\colon \mathbb{P}_i^+ \to \mathbb{P}_i$ i exact in all rectangles.

Let $\{p_\alpha : \alpha<\omega_1\} \subseteq \bigcup\{\mathbb{P}_i : i<\delta\}$ be a maximal antichain. We show that it is also a maximal antichain in $\bigcup\{\mathbb{P}_i^+ : i<\delta\}$. Suppose $p \in \mathbb{P}_i^+$ is incompatible with all the p_α's. But then $f_{i^+,i}(p)$ is incompatible in $\bigcup\{\mathbb{P}_i : i<\delta\}$ with the alledged maximal antichain.

Now we have to show that also the rectangles with limits in the right hand corners are exact: Let $\mathbb{P} = \bigcup_{\alpha<\delta}\mathbb{P}_\alpha$ and $\mathbb{P}^+ = \bigcup_{\alpha<\delta}\mathbb{P}_\alpha^+$ and let $i<\delta$, $p_i^+ \in \mathbb{P}_i^+$. Then

$$\begin{aligned}
&\sup_{\mathbb{P}_i}\{p_i \in \mathbb{P}_1 : p_i \perp_{\mathbb{P}_i^+} p_i^+\}\\
&= \sup_{\mathbb{P}_\alpha}\{p_\alpha \in \mathbb{P}_\alpha : p_\alpha \perp_{\mathbb{P}_\alpha^+} p_1^+\}\\
&= \sup_{\mathbb{P}_\delta}\{p_\delta \in \mathbb{P}_\delta : p_\delta \perp_{\mathbb{P}_\delta^+} p_1^+\},
\end{aligned}$$

because the rectangle from i to α is exact and because $\mathbb{P}_\delta^+$ is the direct limit of the P_α's. But are the *limit* triples $(\bigcup_{\alpha<\delta}\mathbb{P}_\alpha, \bigcup_{\alpha<\delta}\mathbb{P}_\alpha^+, \underset{\sim}{\eta})$ in Γ? Say we formulate it for Γ_{new} and the proof of this item of Definition 2.1 can be used also for Γ_{Cohen}, Γ_{ud}, Γ_{tow}, $\Gamma_{\mathbf{q}}$ (from Claim 2.5).

The essential part is to show (for (a) for Γ_{new}) that

$$\Vdash_{\bigcup\{\mathbb{P}_i^+ : i<\delta\}} \underset{\sim}{\eta} \notin \mathbf{V}^{\bigcup\{\mathbb{P}_i : i<\delta\}}.$$

For this we use "Tools for your forcing construction" [6] p. 358, the part for the c.c.c. forcings and the finite support iteration (taking unions of increasing notions of forcings is the same as finite support iteration). Again we can use common properties of Γ_{new}, Γ_{Cohen}, Γ_{ud}, Γ_{tow}, $\Gamma_{\mathbf{q}}$.

We write $\bigcup\{\mathbb{P}_i^+ : i<\delta\} = \mathbb{P}_\delta^+$. $\bigcup\{\mathbb{P}_i : i<\delta\} = \mathbb{P}_\delta$.

Note that all the time, for all our five Γ's, for the limit property, we use the same fact:

$$\Vdash_{\mathbb{P}^+} \text{"}\underset{\sim}{\eta} \text{ is in } \Gamma\text{-relation with all the reals of } V^{\mathbb{P}}\text{"}$$

is of the form

$$\Vdash_{\mathbb{P}^+} (\forall \nu \in \omega^\omega \cap V^{\mathbb{P}}) \neg \underset{\sim}{\eta} R \nu,$$

where $R = R_\Gamma$ is an F_σ relation. There is a preservation theorem for this situation, a modification of Theorem 8.4 of [6].

Hypothesis 2.6. *We assume that* $R = \sqsubset = \bigcup\{\sqsubset_n : n < \omega\}$ *and for all* n *and* $\eta \in \mathbb{R}$ *the set* $\{g \in \mathbf{V}^{\mathbb{P}} : \eta \sqsubset_n g\}$ *is closed.*

This is true for $R = R_\Gamma$ being the equality for item (a), being $\leq^* = \bigcup\{\{(f,g) : f \restriction [n,\infty) \leq g \restriction [n,\infty)\} : n < \omega\}$ for item (b), being $\sqsubset_{\text{Cohen}}$ (see [2, Chapter 7]) for item (c) and $\leq_{\text{tow}} = \bigcup \leq_{\text{tow},n}$ with $\eta \leq_{\text{tow},n} g$ iff $\eta^{-1}\{1\} \cap g^{-1}\{1\} \cap [n,\infty) = \emptyset$.

Theorem 2.7. *(Modification of Theorem 8.4 of* [6]*). Let* $\langle \mathbb{P}^+_\alpha : \alpha < \delta \rangle$ *be a* $\lessdot$ *increasing continuous sequence of c.c.c. forcings and let* $\mathbb{P}_\delta$ *be their direct limit. Let* $\langle \mathbb{P}_\alpha, : \alpha < \delta \rangle$ *be a sequence such that all the rectangles from the diagramme* (2.3) *are exact. Let* $\underset{\sim}{\eta}$ *be a* $\mathbb{P}^+_0$*-name such that*

$$\forall \alpha < \delta \Vdash_{\mathbb{P}^+_\alpha} \forall g \in \mathbf{V}^{\mathbb{P}_\alpha} (\underset{\sim}{\eta} \not\sqsubset g).$$

Then

$$\Vdash_{\mathbb{P}^+_\delta} \forall g \in \mathbf{V}^{\mathbb{P}_\delta} (\underset{\sim}{\eta} \not\sqsubset g).$$

Proof. Let $\langle \delta_i : i < \text{cf}(\delta) \rangle$ be sequence of ordinals converging to δ. We assume that $\text{cf}(\delta) = \omega$, because otherwise there is nothing to show because by the c.c.c. each real will appear for the first time at a stage with cofinality less of equal ω. Assume the conclusion is false, so there is a condition $p_0 \in \mathbb{P}^+_\delta$ and a $\mathbb{P}_\delta$-name $\underset{\sim}{g}$ such that $p_0 \Vdash_{\mathbb{P}^+_\delta} \underset{\sim}{\eta} \sqsubset \underset{\sim}{g}$. We find a condition $p_1 \leq p_0$ and an integer n such that $p_1 \Vdash \underset{\sim}{\eta} \sqsubset_n \underset{\sim}{g}$. Since p_1 has finite support there is some $i < \text{cf}(\delta)$ such that $p_1 \in \mathbb{P}^+_{\delta_i}$.

Let G be $\mathbb{P}^+_\delta$ -generic over $\mathbf{V}$, and let G_{δ_i} be $G \cap \mathbb{P}_{\delta_i}$. Clearly in $\mathbf{V}[G_{\delta_i}]$ with $p_1 \in G_{\delta_i}$ we have that

$$\Vdash_{\delta_i,\delta} \underset{\sim}{\eta} \sqsubset_n \underset{\sim}{g}. \tag{2.4}$$

On the other hand $\underset{\sim}{\eta}[G_{\delta_i}] \not\sqsubset_n x$ for all $x \in V[G_{\delta_i}]$. Let $\text{tree}(\underset{\sim}{g}[G_{\delta_i}])$ be the tree of possible evaluations of $\underset{\sim}{g}[G_{\delta_i}]$, all possible evaluations, not only generic ones. Take $x \in (\text{tree}(\underset{\sim}{g}[G_{\delta_i}]))^{\mathbf{V}[G_{\delta_i}]}$.

We consider the set $\{h : \underset{\sim}{\eta} \not\sqsubset_n h\} \subseteq \mathbf{V}[G_\delta]$. This is an open set containing x. So there is some $k < \omega$ such that

$$\Vdash_{\mathbb{P}_{\delta_i,\delta}} \forall h \, (h \restriction k = \underset{\sim}{g}[G_{\delta_i}] \restriction k \rightarrow \underset{\sim}{\eta}[G_{\delta_i}] \not\sqsubset_n h).$$

Let $q \in P_\delta/G_{\delta_i}$ be a condition forcing $\underset{\sim}{g} \restriction k = x \restriction k$. Then also q forces in $\mathbb{P}_{\delta_i,\delta}$ that $\underset{\sim}{\eta} \not\sqsubset_n \underset{\sim}{g}$ in contradiction to (2.4). $\square_{\text{limit part of }2.5}$

Now for the *amalgamation*: Let $(\mathbb{P}_1, \mathbb{P}_1^+, \underset{\sim}{\eta}) \in \Gamma$, let $\mathbb{P}_1 \lessdot \mathbb{P}_2$, and let $\mathbb{P}_2/\mathbb{P}_1$ have the Knaster property. Then choose $f \colon \mathbb{P}_2 \to \mathbb{P}_2'$ such that $f \restriction \mathbb{P}_1 = id$ and $\mathbb{P}_2' \cap \mathbb{P}_1^+ = \mathbb{P}_1$. Since $\mathbb{P}_2/\mathbb{P}_1$ has the Knaster property $\mathbb{P}_2^+ = \mathbb{P}_1^+ \times_{\mathbb{P}_1} \mathbb{P}_2 = \mathbb{P}_1 * (\mathbb{P}_1^+/\mathbb{P}_1 \times \mathbb{P}_2/\mathbb{P}_1)$ has the c.c.c.

Then $\mathbb{P}_1^+ \lessdot \mathbb{P}_2^+$. and $id \colon \mathbb{P}_2 \to \mathbb{P}_2^+$ is a complete embedding over $\mathbb{P}_1$.

Theorem 2.8. *Let $(\mathbb{P}_1, \mathbb{P}_1^+, \underset{\sim}{\eta}) \in \Gamma$, $R_\Gamma = \sqsubset$, let $\mathbb{P}_1 \lessdot \mathbb{P}_2$, and let $\mathbb{P}_2/\mathbb{P}_1$ have the Knaster property. Let $\underset{\sim}{\eta}$ be a $\mathbb{P}_1^+$-name such that*

$$\Vdash_{\mathbb{P}_1^+} \forall g \in \mathbf{V}^{\mathbb{P}_1} (\underset{\sim}{\eta} \not\sqsubset g).$$

Then for $\mathbb{P}_2^+ = \mathbb{P}_1^+ \times_{\mathbb{P}_1} \mathbb{P}_2$ we have that

$$\Vdash_{\mathbb{P}_2^+} \forall g \in \mathbf{V}^{\mathbb{P}_2} (\underset{\sim}{\eta} \not\sqsubset g).$$

Proof. Let $(\mathbb{P}_1, \mathbb{P}_1^+, \underset{\sim}{\eta}) \in \Gamma$, let $\mathbb{P}_1 \lessdot \mathbb{P}_2$, and let $\mathbb{P}_2/\mathbb{P}_1$ have the Knaster property. Then choose an isomorphism $f \colon \mathbb{P}_2 \to \mathbb{P}_2'$ such that $f \restriction \mathbb{P}_1 = id$ and $\mathbb{P}_2' \cap \mathbb{P}_1^+ = \mathbb{P}_1$. Since $\mathbb{P}_2/\mathbb{P}_1$ has the Knaster property $\mathbb{P}_2^+ = \mathbb{P}_1^+ \times_{\mathbb{P}_1} \mathbb{P}_2'$ has the c.c.c.

Then $\mathbb{P}_1^+ \lessdot \mathbb{P}_2^+$ and $f \colon \mathbb{P}_2 \to \mathbb{P}_2^+$ is a complete embedding over $\mathbb{P}_1$. Then $\Vdash_{\mathbb{P}_2^+}$ $(\forall g \in \mathbf{V}^{\mathbb{P}_2} \bigwedge_{n<\omega} \eta \not\sqsubset_n g)$ since otherwise there would be some $n < \omega$ and $g \in \mathbf{V}^{\mathbb{P}_2}$ and p_2^+ such that $p_2^+ \Vdash_{\mathbb{P}_2^+} \eta \sqsubset_n g$. Let G_2^+ be $\mathbb{P}_2^+$-generic over V and let $p_2^+ \in G_2^+$ and $G_2 = G_2^+ \cap \mathbb{P}_2$. Then $T = \{s \in \omega^{<\omega} \,:\, (\exists g \in V^{\mathbb{P}_2})(s \subseteq g \wedge \eta[G_2^+] \sqsubset_n g)\}$ is an ill-founded tree in $V[G_2^+]$. But since $\sqsubset_n$ is absolute and since $\eta[G_2^+] = \eta[G_2^+ \cap \mathbb{P}_1^+]$ we have that the T is ill-founded is $V[G_2^+ \cap \mathbb{P}_1^+]$ as well and $G_2^+ \cap \mathbb{P}_1^+$ is $\mathbb{P}_1^+$-generic over $\mathbf{V}$, and this is a contradiction. $\square_{\text{amalgamation part of }2.5}$

Now for the LS property: Since it has the c.c.c., $\underset{\sim}{\eta}$ is determined by countably many countable antichains.

$|Y| = \mu < \lambda$, and hence by closing $\mathbb{P} \cap Y$ under the suprema in $\mathbb{P}$ of all subsets of $\mathbb{P}$, we get $\mathbb{P}_1 \lessdot \mathbb{P}$ such that $\mathbb{P} \cap Y \subseteq \mathbb{P}_1$. We can require that $|\mathbb{P}_1| < \lambda$, because every supremum is determined by countably many elements and hence there are less or equal than $\mu^{\aleph_0} < \lambda$ many tasks. We do the same for Y in $\mathbb{P}^+$ and for $\mathbb{P}_1$ and thus find $\mathbb{P}_1^+$. Then we automatically have $\mathbb{P}_1 \lessdot \mathbb{P}_1^+$ and $\mathbb{P} \times_{\mathbb{P}_1} \mathbb{P}_1^+ \lessdot \mathbb{P}^+$, and thus get the diagramme

$$\begin{tikzcd}
\mathbb{P} \arrow[r, "\lessdot"] & \mathbb{P}^+ \\
\mathbb{P}_1 \arrow[r, "\lessdot"] \arrow[u, "\lessdot"] & \mathbb{P}_1^+ \arrow[u, "\lessdot"'] \\
\mathbb{P} \cap Y \arrow[r, "\subseteq"] \arrow[u, "\subseteq"] & Y \arrow[u, "\subseteq"']
\end{tikzcd}$$

whose upper square is exact. This was the proof for Γ_{new} and all other Γ's, because it did not use any specific property but Hypothesis 2.6 $\square_{2.5}$

Definition 2.9. *Let* $\mathbf{q}$ *be a* **forgetful** *pcrf, and let* Γ *be a forcing bigness notion.*

1. *We say "*$\mathbf{q}$ *is orthogonal to* Γ*" and write* $\mathbf{q} \perp \Gamma$*, if the following holds: If*
   - *(a)* $(\mathbb{P}_1, \mathbb{P}_1^+, \underset{\sim}{\eta}) \in \Gamma$ *and*
   - *(b)* $\mathbb{P}_1 \lessdot \mathbb{P}_2$ *and* $\mathbb{P}_2/\mathbb{P}_1$ *is Cohen*$_A$ *for some set* A *and* $\mathbb{P}_2 \cap \mathbb{P}_1^+ = \mathbb{P}_1$,
   - *(c)* $\underset{\sim}{\mathcal{C}}_2$ *is a* $\mathbb{P}_2$*-name and* $\mathbb{P}_2$ *forces that it is* $\mathbf{q}$*-nice,*

   then we can find $(\mathbb{P}_2^+, f, \underset{\sim}{\mathcal{C}}_2^+)$ *such that*
   - *(α)* $\mathbb{P}_1^+ \lessdot \mathbb{P}_2^+$,
   - *(β)* f *is a complete embedding of* $\mathbb{P}_2$ *into* $\mathbb{P}_2^+$ *over* $\mathbb{P}_1$,
   - *(γ)* $\Vdash_{\mathbb{P}_2^+}$ "$f(\underset{\sim}{\mathcal{C}}_2) \subseteq \underset{\sim}{\mathcal{C}}_2{}^+ \wedge \underset{\sim}{\mathcal{C}}_2{}^+$ *is* $\mathbf{q}$*-nice*",
   - *(δ)* $(\mathbb{P}_1, \mathbb{P}_1^+, \underset{\sim}{\eta}) \leq_\Gamma (f(\mathbb{P}_2), \mathbb{P}_2^+, \underset{\sim}{\eta}) \leq_\Gamma (f(\mathbb{P}_2) * \mathbb{A}(f(\underset{\sim}{\mathcal{C}}_2)), \mathbb{P}_2^+ * \mathbb{A}(\underset{\sim}{\mathcal{C}}_2{}^+), \underset{\sim}{\eta})$.

   We say that $\mathbf{q} \perp \Gamma$ *with a simple witness, if in the conclusion we can choose* $\mathbb{P}_2^+ = (\mathbb{P}_1^+ \times_{\mathbb{P}_1} \mathbb{P}_2) \times \text{Cohen}_\lambda$, $\lambda = (|\mathbb{P}_1^+| + |\mathbb{P}_2|)^{\aleph_0}$, $f = id$.

2. *We say that* $(\mathbb{P}, \mathbb{P}^+, \underset{\sim}{\eta}, \underset{\sim}{\mathcal{C}}, \underset{\sim}{\mathcal{C}}^+)$ *is a* $(\mathbf{q}, \Gamma)$*-tuple if*
   - *(a)* $(\mathbb{P}, \mathbb{P}^+, \underset{\sim}{\eta}) \in \Gamma$,
   - *(b)* $\underset{\sim}{\mathcal{C}}$ *is a* $\mathbb{P}$*-name and* $\mathbb{P}$ *forces that it is* $\mathbf{q}$*-nice,*
   - *(c)* $\underset{\sim}{\mathcal{C}}^+$ *is a* $\mathbb{P}^+$*-name and* $\mathbb{P}^+$ *forces that it is* $\mathbf{q}$*-nice,*
   - *(d)* $\underset{\sim}{\mathcal{C}} = \underset{\sim}{\mathcal{C}}^+ \cap \mathbf{Q}^{\mathbf{V}[\mathbb{P}]}$
   - *(e)* $(\mathbb{P}, \mathbb{P}^+, \underset{\sim}{\eta}) \leq_\Gamma (\mathbb{P} * \mathbb{A}(\underset{\sim}{\mathcal{C}}), \mathbb{P}^+ * \mathbb{A}(\underset{\sim}{\mathcal{C}}^+), \underset{\sim}{\eta})$.

3. *We say "$\mathbf{q}$ is strongly orthogonal to Γ" and write $\mathbf{q} \perp_{st} \Gamma$, if: If*
   (a) *Γ is a forcing bigness notion,*
   (b) *$\mathbf{q}$ is a forgetful pcrf,*
   (c) *$(\mathbb{P}_1, \mathbb{P}_1^+, \underset{\sim}{\eta}, \underset{\sim}{\mathcal{C}}_1, \underset{\sim}{\mathcal{C}}_1^+)$ is a $(\mathbf{q}, \Gamma)$-tuple and if $\mathbb{P}_1 \lessdot \mathbb{P}_2$ and $\mathbb{P}_2/\mathbb{P}_1$ satisfies the Knaster condition and $\Vdash_{\mathbb{P}_2} \underset{\sim}{\mathcal{C}}_2$ is a $\mathbf{q}$-nice family extending $\underset{\sim}{\mathcal{C}}_1$,*

   then we can find $(\mathbb{P}_2^+, f, \underset{\sim}{\mathcal{C}}_2^+)$ such that
   (α) *$\mathbb{P}_1^+ \lessdot \mathbb{P}_2^+$,*
   (β) *f is a complete embedding of $\mathbb{P}_2$ into $\mathbb{P}_2^+$ over $\mathbb{P}_1$,*
   (γ) *$\Vdash_{\mathbb{P}_2^+}$ "$f(\underset{\sim}{\mathcal{C}}_2) \subseteq \underset{\sim}{\mathcal{C}}_2^+$",*
   (δ) *$(\mathbb{P}_2, \mathbb{P}_2^+, \underset{\sim}{\eta}, f(\underset{\sim}{\mathcal{C}}_2), {\underset{\sim}{\mathcal{C}}_2}^+)$ is a $(\mathbf{q}, \Gamma)$-tuple,*
   (ε) *$(\mathbb{P}_1 * \mathbb{A}(\underset{\sim}{\mathcal{C}}_1), \mathbb{P}_1^+ * \mathbb{A}(\underset{\sim}{\mathcal{C}}_1^+), \underset{\sim}{\eta}) \leq_\Gamma (\mathbb{P}_2 * \mathbb{A}(f(\underset{\sim}{\mathcal{C}}_2)), \mathbb{P}_2^+ * \mathbb{A}(\underset{\sim}{\mathcal{C}}_2^+), \underset{\sim}{\eta})$.*

4. *The order $\leq$ on the family of $(\mathbf{q}, \Gamma)$-tuples is defined by*

$$(\mathbb{P}_1, \mathbb{P}_1^+, \underset{\sim}{\eta}, \underset{\sim}{\mathcal{C}}_1, \underset{\sim}{\mathcal{C}}_1^+) \leq (\mathbb{P}_2, \mathbb{P}_2^+, \underset{\sim}{\eta}, \underset{\sim}{\mathcal{C}}_2, \underset{\sim}{\mathcal{C}}_2^+)$$

   if both are $(\mathbf{q}, \Gamma)$-tuples and
   (α) *$\mathbb{P}_1 \lessdot \mathbb{P}_2$,*
   (β) *$\mathbb{P}_1^+ \lessdot \mathbb{P}_2^+$,*
   (γ) *$\underset{\sim}{\eta}_1 = \underset{\sim}{\eta}_2$,*
   (δ) *$\Vdash_{\mathbb{P}_2}$ "$\underset{\sim}{\mathcal{C}}_1 \subseteq \underset{\sim}{\mathcal{C}}_2$",*
   (ε) *$\Vdash_{\mathbb{P}_2^+}$ "$\underset{\sim}{\mathcal{C}}_1^+ \subseteq \underset{\sim}{\mathcal{C}}_2^+$".*
   (ζ) *$(\mathbb{P}_1 * \mathbb{A}(\underset{\sim}{\mathcal{C}}_1), \mathbb{P}_1^+ * \mathbb{A}(\underset{\sim}{\mathcal{C}}_1^+), \underset{\sim}{\eta}) \leq_\Gamma (\mathbb{P}_2 * \mathbb{A}(f(\underset{\sim}{\mathcal{C}}_2)), \mathbb{P}_2^+ * \mathbb{A}(\underset{\sim}{\mathcal{C}}_2^+), \underset{\sim}{\eta})$.*

Claim 2.10. *Assume that Γ is a c.c.c. forcing bigness notion with amalgamation. A sufficient condition for $\mathbf{q} \perp \Gamma$ is the following:*

⊠ *If*

(a) *$(\mathbb{P}, \mathbb{P}^+, \underset{\sim}{\eta}) \in \Gamma$,*

(b) *$\Vdash_{\mathbb{P}}$ "$\underset{\sim}{\mathfrak{c}} \in \mathbf{Q}$" or just for some $(\mathbb{P}_0, \underset{\sim}{\mathcal{C}})$ we have that $\mathbb{P} \lessdot \mathbb{P}_0$ and $\Vdash_{\mathbb{P}_0}$ "$\underset{\sim}{\mathcal{C}}$ is $\mathbf{q}$-nice" and $\Vdash_{\mathbb{P}_0}$ "$\underset{\sim}{\mathfrak{c}}$ is generic for $(\mathbf{Q}(\underset{\sim}{\mathcal{C}}))^{\mathbf{V}[G_{\mathbb{P}}]}$",*

then we can find $\underset{\sim}{\mathcal{C}}^+$ and a $\mathbb{P}^+$-name of a $\mathbf{q}$-nice family to which $\bar{\mathfrak{c}}$ belongs such that if $\mathcal{J}$ is a $\mathbb{P}$-name of a subset of $\mathbb{A}_{\mathcal{C}}(\bar{\mathfrak{c}})$ which is explicitly predense then $\Vdash_{\mathbb{P}^+}$ "$\mathcal{J}$ is a predense subset of $\mathbb{A}(\mathcal{C}^+)$" and $(\mathbb{P} * \mathbb{A}(\underset{\sim}{\mathcal{C}}), \mathbb{P}^+ * \mathbb{A}(\underset{\sim}{\mathcal{C}}^+), \underset{\sim}{\eta}) \in \Gamma$.

Proof. Suppose $\mathbb{P}_1$, $\mathbb{P}_1^+$, $\underset{\sim}{\eta}$, $\mathbb{P}_2$ and $\underset{\sim}{\mathcal{C}}_2$ are given as in the premises (a), (b) and (c) of Definition 2.9 Part 1. We have to find $(\mathbb{P}_2^+, f, \underset{\sim}{\mathcal{C}}_2^+)$ as there.

Since Γ has the amalgamation property, there is $(\mathbb{P}_1, \mathbb{P}_1^+, \underset{\sim}{\eta}) \leq_\Gamma (f(\mathbb{P}_2), \mathbb{P}_2^+, \underset{\sim}{\eta})$. Now we read the condition given in Claim 2.10 for $(f(\mathbb{P}_2), \mathbb{P}_2^+, \underset{\sim}{\eta})$ and $\underset{\sim}{\mathcal{C}}_2$ and find $\underset{\sim}{\mathcal{C}}_2^+$ such that $(\mathbb{P}_1, \mathbb{P}_1^+, \underset{\sim}{\eta}) \leq_\Gamma (f(\mathbb{P}_2), \mathbb{P}_2^+, \underset{\sim}{\eta}) \leq_\Gamma (f(\mathbb{P}_2) * \mathbb{A}(f(\underset{\sim}{\mathcal{C}}_2)), \mathbb{P}_2^+ * \mathbb{A}(\underset{\sim}{\mathcal{C}}_2^+), \underset{\sim}{\eta})$. □

3. LONG LOW C.C.C. ITERATION AND $\mathbf{q} \perp \Gamma$

Hypothesis 3.1. $\lambda = \mathrm{cf}(\lambda)$ *satifies*

$$(\forall \alpha < \lambda)(|\alpha|^{\aleph_0} < \lambda),$$

and $\zeta(*)$ is a limit of uncountable cofinality.

Definition 3.2. *Let $\bar{\Gamma} = \langle \Gamma_\zeta : \zeta < \zeta(*) \rangle$ be sequence of c.c.c. forcing bigness notions.*

- *(0)* *$\bar{\Gamma}$ is nice if each Γ_ζ is, i.e. Γ_ζ is a c.c.c. forcing bigness notion, has amalgamation and the $(< \lambda)$-L.S. $\bar{\Gamma}$ has free amalgamation if each Γ_ζ has.*
- *(1)* *$K = K_{\lambda, \zeta(*)}(\bar{\Gamma})$ is the family of $\mathfrak{q} = \langle \mathbb{P}_\zeta, \underset{\sim}{\eta}_\zeta : \zeta < \zeta(*) \rangle = \langle \mathbb{P}_\zeta^{\mathfrak{q}}, \underset{\sim}{\eta}_\zeta^{\mathfrak{q}} : \zeta < \zeta(*) \rangle$ such that*
  - *(a)* *$\mathbb{P}_\zeta$ is a c.c.c. forcing notion of cardinality $< \lambda$ for $\zeta \leq \zeta(*)$.*
  - *(b)* *$\langle \mathbb{P}_\zeta : \zeta < \zeta(*) \rangle$ is $\lessdot$-increasing, and if $\xi < \zeta(*)$ is a limit ordinal, then $\mathbb{P}_{<\xi} = \bigcup \{\mathbb{P}_\zeta : \zeta < \xi\} \lessdot \mathbb{P}_\xi$, if $\aleph_0 < \mathrm{cf}(\xi)$.*
  - *(c)* *$\underset{\sim}{\eta}_\zeta$ is a $\mathbb{P}_{\zeta+1}$ name such that $(\mathbb{P}_\zeta, \mathbb{P}_{\zeta+1}, \underset{\sim}{\eta}_\zeta)$ is Γ_ζ-big.*

  We stipulate $\mathbb{P}_{\zeta(*)}^{\mathfrak{q}} = \bigcup \{\mathbb{P}_\zeta^{\mathfrak{q}} : \zeta < \zeta(*)\}$ and we may write $\mathbb{P}_\zeta[\mathfrak{q}]$ instead of $\mathbb{P}_\zeta^{\mathfrak{q}}$.
- *(1a)* *We say $\mathfrak{q}$ is λ-standard if $\mathfrak{q} \in \mathcal{H}(\lambda)$ and we write $K^* = K_{\lambda, \zeta(*)}^*(\bar{\Gamma}) := K_{\lambda, \zeta(*)}(\bar{\Gamma}) \cap \mathcal{H}(\lambda)$ when $\bar{\Gamma}$, λ, $\zeta(*)$ are clear from the context.*
- *(2)* *We define three binary relations $\leq$, $\leq_{\mathrm{pr}}$ and $\leq_{ap}$ on K^*:*
  - *(a)* *$\mathfrak{p} \leq_{pr} \mathfrak{q}$ iff $\zeta < \zeta(*)$ implies $\underset{\sim}{\eta}_\zeta^{\mathfrak{p}} = \underset{\sim}{\eta}_\zeta^{\mathfrak{q}}$ and $\mathbb{P}_\zeta^{\mathfrak{p}} \lessdot \mathbb{P}_\zeta^{\mathfrak{q}}$,*
  - *(b)* *$\mathfrak{p} \leq \mathfrak{q}$ iff there is some $\xi < \zeta(*)$ such that $\zeta \in [\xi, \zeta(*))$ the relations $\underset{\sim}{\eta}_\zeta^{\mathfrak{p}} = \underset{\sim}{\eta}_\zeta^{\mathfrak{q}}$ and $\mathbb{P}_\zeta^{\mathfrak{p}} \lessdot \mathbb{P}_\zeta^{\mathfrak{q}}$ hold,*
  - *(c)* *$\mathfrak{p} \leq_{ap} \mathfrak{q}$ iff $\mathfrak{p} \leq \mathfrak{q}$ and $(\forall^* \zeta < \zeta(*))(\underset{\sim}{\eta}_\zeta^{\mathfrak{p}} = \underset{\sim}{\eta}_\zeta^{\mathfrak{q}} \wedge \mathbb{P}_\zeta^{\mathfrak{p}} = \mathbb{P}_\zeta^{\mathfrak{q}})$,*
  - *(d)* *$\mathfrak{p} \equiv \mathfrak{q}$ iff $\mathfrak{p} \leq_{ap} \mathfrak{q} \leq_{ap} \mathfrak{p}$.*

(3) *If* $\langle \mathfrak{q}_\alpha : \alpha < \delta \rangle$ *is* $\leq_{pr}$*-increasing, let* $\mathfrak{q}_\delta = \bigcup_{\alpha<\delta} \mathfrak{q}_\alpha$ *be defined as* $\langle \bigcup_{\alpha<\delta} \mathbb{P}^{\mathfrak{q}_\alpha}_\zeta, \underset{\sim}{\eta}_\zeta : \zeta < \zeta(*) \rangle$.

(4) $\mathfrak{K}^*$ *as a forcing notion is* $(K^*, \leq)$ *and the generic is* $\underset{\sim}{\mathbb{P}} = \bigcup \{ \mathbb{P}^{\mathfrak{q}}_{\zeta(*)} : \mathfrak{q} \in \underset{\sim}{G}_{K^*} \}$.

(5) *If for all* $\zeta < \zeta(*)$, $\Gamma_\zeta = \Gamma$, *then we may write* Γ *instead of* $\bar{\Gamma}$.

(6) *We say* $\langle \mathfrak{q}_\alpha : \alpha < \alpha^* \rangle$ *is* $\leq_{pr}$*-continuous if*

  (a) $\alpha < \beta < \alpha^* \Rightarrow \mathfrak{q}_\alpha \leq_{pr} \mathfrak{q}_\beta$.

  (b) *If* $\beta < \alpha^*$ *is a limit ordinal then* $\mathfrak{q}_\beta = \bigcup_{\alpha<\beta} \mathfrak{q}_\alpha$, *i.e.,* $\mathbb{P}^{\mathfrak{q}_\beta}_\zeta = \bigcup \{ \mathbb{P}^{\mathfrak{q}_\alpha}_\zeta : \alpha < \beta \}$ *for every* $\zeta < \zeta(*)$.

(7) *We replace continuous by "weakly continuous" if in clause (b) above only* $\bigcup \{ \mathbb{P}^{\mathfrak{q}_\alpha}_\zeta : \alpha < \beta \} \lessdot \mathbb{P}^{\mathfrak{q}_\beta}_\zeta$ *is demanded.*


$\zeta(*)$
$\langle \mathbb{P}^{\mathfrak{q}}_\zeta, \underset{\sim}{\eta}^{\mathfrak{q}}_\zeta : \zeta < \zeta(*) \rangle$
$(\mathbb{P}_\zeta, \mathbb{P}_{\zeta+1}, \underset{\sim}{\eta}) \in \Gamma_\zeta$
Γ_ζ
$\mathfrak{p}$
$\mathfrak{q}$
λ

Figure 1: A sketch of $K_{\lambda,\zeta(*)}(\bar{\Gamma})$

Claim 3.3. (1) $\leq$ *and* $\leq_{pr}$ *are partial orders on* K^*.

(2) $\equiv$ *is an equivalence relation on* K^* *and the forcing* $(K^*, \leq)$ *is equivalent to* $(K^*/\equiv, \leq)$.

(3) *If* Γ *is nice and if* $\delta < \lambda$ *is a limit ordinal and* $\langle \mathfrak{q}_\alpha : \alpha < \delta \rangle$ *is* $\leq_{pr}$*-increasing and continuous as a sequence of members in* K^* *and* $\mathfrak{q}_\delta = \bigcup_{\alpha<\delta} \mathfrak{q}_\alpha$ *then* $\mathfrak{q}_\delta \in K^*$ *and* $\langle \mathfrak{q}_\alpha : \alpha \leq \delta \rangle$ *is* $\leq_{pr}$*-increasing and continuous. Instead of continuous we can demand weakly continuous.*

(4) $\mathfrak{p} \equiv \mathfrak{q} \Rightarrow \mathfrak{p} \leq \mathfrak{q}$ *and* $\mathfrak{p} \leq_{pr} \mathfrak{q} \Rightarrow \mathfrak{p} \leq \mathfrak{q}$.

(5) *As each* Γ_ζ *has the* $(<\lambda)$*-L.S., clearly* $K^*_{\lambda,\zeta(*)} \neq \emptyset$.

(6) *If each Γ_ζ, $\zeta < \zeta(*)$, has free amalgamation and $\mathfrak{q} \in K^*_{\lambda,\zeta(*)}$ and $\xi < \zeta(*)$ and $\mathbb{P}^{\mathfrak{q}}_\xi \lessdot \mathbb{P} \in \mathcal{H}(\lambda)$ and $\mathbb{P}$ is c.c.c. and $\Vdash_{\mathbb{P}^{\mathfrak{q}}_{\zeta(*)}}$ "$\mathbb{P}/(\underset{\sim}{G} \cap \mathbb{P}^{\mathfrak{q}}_\xi)$ satisfies the c.c.c.".* then we can find some $(\mathfrak{q}^+, f)$ such that

- (a) $\mathfrak{q} \le_{pr} \mathfrak{q}^+ \in K^*$,
- (b) $\zeta < \xi \Rightarrow \mathbb{P}^{\mathfrak{q}^+}_\zeta = \mathbb{P}^{\mathfrak{q}}_\zeta$
- (c) f is a $\lessdot$-complete embedding of $\mathbb{P}$ into $\mathbb{P}^{\mathfrak{q}^+}_\xi$ over $\mathbb{P}^{\mathfrak{q}}_\xi$:

$$\begin{tikzcd}
\mathbb{P}^{\mathfrak{q}}_{\xi+1} \arrow[rr, "\lessdot"] & & \mathbb{P}^{\mathfrak{q}^+}_{\xi+1} \\
\mathbb{P}^{\mathfrak{q}}_{\xi} \arrow[u, "\lessdot"] \arrow[r, "\lessdot"] & \mathbb{P} \arrow[r, "{\lessdot, f}"] & \mathbb{P}^{\mathfrak{q}^+}_{\xi} \arrow[u, "\lessdot"]
\end{tikzcd}$$

Proof. This follows almost immediately from the definitions. □

Claim 3.4. *(1) If $\mathfrak{p} \le \mathfrak{q}$ then for some $\mathfrak{q}' \equiv \mathfrak{q}$ we have $\mathfrak{p} \le_{pr} \mathfrak{q}'$.*

(2) *If $\underset{\sim}{G}_{K^*} \subseteq K^*_{\lambda,\zeta(*)}$ is generic over $\mathbf{V}$, then $\underset{\sim}{\mathbb{P}} = \bigcup\{\mathbb{P}^{\mathfrak{p}}_{\zeta(*)} : \mathfrak{p} \in \underset{\sim}{G}_{K^*}\}$ is a c.c.c. forcing notion of cardinality $\le \lambda$ and $\mathfrak{q} \Vdash \mathbb{P}^{\mathfrak{q}}_{\zeta(*)} \lessdot \underset{\sim}{\mathbb{P}}$, but it locally looks like $\mathbb{P}^{\mathfrak{q}}_{\zeta(*)}$.*

(3) *If $\bar{\Gamma}$ is nice and if $\mathfrak{p} \in K^*$ and if $\xi < \zeta(*)$ and if we define $\mathfrak{q}$ by*

$$\mathbb{P}^{\mathfrak{q}}_\zeta = \begin{cases} \mathbb{P}^{\mathfrak{p}}_\zeta & \text{if } \zeta < \xi, \\ \mathbb{P}^{\mathfrak{p}}_\zeta * ({}^{\omega>}\omega, \lhd) & \text{if } \zeta \in [\xi, \zeta(*)), \end{cases}$$

then $\mathfrak{p} \le_{pr} \mathfrak{q} \in K^*$.

Claim 3.5. *Let $G_{\mathfrak{K}^*} \subseteq \mathfrak{K}^*$ be generic over $\mathbf{V}$ and let $\mathbb{P} = \underset{\sim}{\mathbb{P}}[G_{\mathfrak{K}^*}]$. Then in $\mathbf{V}[G_{\mathfrak{K}}]$:*

(a) *$\mathbb{P}$ is a c.c.c. forcing notion of cardinality λ adding λ reals.*

(b) *If $\underset{\sim}{\nu}$ is a $\mathbb{P}$-name for a real and $\mathfrak{q}^* = \langle \mathbb{P}^*_\zeta, \underset{\sim}{\eta}_\zeta : \zeta < \zeta(*)\rangle \in G_{\mathfrak{K}^*}$, then for every sufficiently large $\zeta < \zeta(*)$ for some $\mathbb{P}_\zeta \lessdot \mathbb{P}^+_\zeta \lessdot \mathbb{P}$ we have*

- *(α) $\underset{\sim}{\nu}$ is a $\mathbb{P}_\zeta$-name,*
- *(β) $(\mathbb{P}_\zeta, \mathbb{P}^+_\zeta, \underset{\sim}{\eta}_\zeta) \in \Gamma_\zeta$.*

(b') *So, if $\zeta < \zeta(*) \Rightarrow \Gamma_\zeta = \Gamma_{ud}$ and $\mathfrak{q}^*$ is as above, then $\mathfrak{q}^* \Vdash_{\mathbb{P}}$ "$\{\underset{\sim}{\eta}_\zeta : \zeta < \zeta(*)\}$ is unbounded" (hence $\mathfrak{b} \le \zeta(*)$).*

(c) *Similar for other relations on ${}^\omega 2$.*

Proof. Item (b): Since $\underset{\sim}{\mathbb{P}}$ has the c.c.c., for $m < \omega$, let $\{p_{m,n} : n < \omega\} \subseteq \mathbb{P}$ be a maximal antichain forcing the value of $\underset{\sim}{\nu} \restriction m$. $\mathfrak{K}^*$ is λ-complete and $\lambda > \aleph_1$, hence there is some $\mathfrak{q} \in G_K$ such that $\{p_{m,n} : n, m \in \omega\} \subseteq \mathbb{P}^{\mathfrak{q}}$ and $\mathfrak{q} \geq \mathfrak{q}^*$.

Since $\mathbb{P}^{\mathfrak{q}}$ is the union of the increasing chain $\langle \mathbb{P}^{\mathfrak{q}}_\zeta : \zeta < \zeta(*)\rangle$ and since $\mathrm{cf}(\zeta(*)) > \aleph_0$, for some $\zeta_0 < \zeta(*)$ we have that $\{p_{m,n} : n, m \in \omega\} \subseteq \underset{\sim}{\mathbb{P}}^{\mathfrak{q}}_{\zeta_0}$.

As $\mathfrak{q}^* \in G$, $\mathfrak{q}^* \leq \mathfrak{q}$ we have for some $\zeta_1 < \zeta(*)$ that $\bigwedge_{\xi \in [\zeta_1, \zeta(*))} (\mathbb{P}^{\mathfrak{q}^*}_\xi \lessdot \mathbb{P}^{\mathfrak{q}}_\xi \wedge \underset{\sim}{\eta}^{\mathfrak{q}^*}_\xi = \underset{\sim}{\eta}^{\mathfrak{q}}_\xi)$.

So let $\zeta \in [\max(\zeta_0, \zeta_1), \zeta(*))$. Since $\mathfrak{q} \in G_{K^*}$, we have $\mathbb{P}^{\mathfrak{q}}_\zeta \lessdot \mathbb{P}^{\mathfrak{q}}_{\zeta+1} \lessdot \underset{\sim}{\mathbb{P}}[G_{K^*}]$. Now since $\underset{\sim}{\nu}$ is a $\mathbb{P}^{\mathfrak{q}}_\zeta$-name, by the demands on $\mathfrak{q}$ all follows. □

Definition 3.6. *Let* $\mathfrak{q}_* \in \mathfrak{K}^*$, $\zeta_* < \zeta(*)$ *and* $\zeta_* + \zeta(*) = \zeta(*)$, *for notational simplicity, and* $p_* \in \mathbb{P}^{\mathfrak{q}_*}_\zeta$.

Define $K^*_{\geq \mathfrak{q}_*} = \{\mathfrak{q} \in K : \mathfrak{q}_* \leq \mathfrak{q}\}$ *and define the function* F *with domain* $K^*_{\geq \mathfrak{q}_*}$ *as follows: if* $\mathfrak{q} \in K^*_{\mathfrak{q}_*}$ *then* $F(\mathfrak{q}) = \langle \mathbb{P}'_\zeta : \zeta < \zeta(*)\rangle$ *where* $\mathbb{P}'_\zeta = \mathbb{P}^*[\mathfrak{q}_{\zeta_*+\zeta}] \restriction \{p \in \mathbb{P}^*[\mathfrak{q}_{\zeta_*+\zeta}] : \mathbb{P}^*[\mathfrak{q}_{\zeta_*+\zeta}] \models p_* \leq p\}$ *and lastly* $K^*_{\mathfrak{q}_*, \zeta_*, p_*} = \mathrm{range}(F)$.

Claim 3.7. *1.* $K^*_{\geq \mathfrak{q}_*}, K^*_{\mathfrak{q}_*, \zeta_*, p_*} \subseteq K^*$.

2. F *preserves the order* $\leq, \leq_{pr}, \leq_{ap}$ *and their negations and the forcing* K^*_0 *above* $(\mathfrak{q}_*, p_*)$ *is the same as the forcing* $K^*_{\mathfrak{q}_*, \zeta_*, p_*}$.

Claim 3.8. *Let* $\mathbf{q} \perp \bar{\Gamma}$. *Let each* Γ_ζ *have free amalgamation. Let* $\mathfrak{K}^* = (K^*_{\lambda, \zeta(*)}(\Gamma), \leq)$. *Then* $\mathfrak{K}^* * \underset{\sim}{\mathbb{P}}$ *is a* $(< \lambda)$*-strategically complete forcing of cardinality* $\leq \lambda^{<\lambda}$ *and in* $\mathbf{V}^{\mathfrak{K}^* * \underset{\sim}{\mathbb{P}}}$, *in the game* $\Game_\kappa(\mathbf{q})$ *the empty player does not have a winning strategy.*

Proof. Assume towards a contradiction that we have for some $\kappa = \mathrm{cf}(\kappa) \in (\aleph_0, \lambda)$ and some $\underset{\sim}{St}$ that $\mathfrak{q} \Vdash_{\mathfrak{K}^*} p \Vdash_{\underset{\sim}{\mathbb{P}}}$ "$\underset{\sim}{St}$ is a name for a winning strategy for the empty player in the game $\Game_\kappa(\mathbf{q})$".

Since $\mathbb{P}$ may be replaced by $\mathbb{P}_{\geq p}$, we may assume $p = 1$. We now choose by induction on $\alpha < \kappa$ a triple $(\mathfrak{q}_\alpha, \underset{\sim}{\mathfrak{c}}_\alpha, \underset{\sim}{\mathfrak{d}}_\alpha)$ satisfying

(t_1) $\mathfrak{q}_\alpha \in K^*$ is $\leq_{pr}$-increasing weakly continuous (in K^*),

(t_2) $\mathfrak{q}_0 = \mathfrak{q}$,

(t_3) $\underset{\sim}{\mathfrak{c}}_\alpha$ is a $\mathbb{P}^{\mathfrak{q}_\alpha}_0$-name and $\underset{\sim}{\mathfrak{d}}_\alpha$ is a $\mathbb{P}^{\mathfrak{q}_\alpha}_0$-name (not just $\mathbb{P}^{\mathfrak{q}_\alpha}_\varepsilon$-names for some ε),

(t_4) $\mathfrak{q}_\alpha \Vdash_{\mathfrak{K}^*} 1 \Vdash_{\underset{\sim}{\mathbb{P}}}$ "$\langle \underset{\sim}{\mathfrak{c}}_\beta, \underset{\sim}{\mathfrak{d}}_\beta : \beta < \alpha\rangle$ is a play of $\Game_\kappa(\mathbf{q})$ in which the empty player uses the strategy $\underset{\sim}{St}$".

Assume that we arrive at stage $\alpha < \kappa$. First we let $\mathfrak{p}_{\alpha,0}$ be $\mathfrak{q}$ if $\zeta = 0$, and we let $\mathfrak{p}_{\alpha,0}$ be $\mathfrak{q}_{\alpha-1}$ if α is a successor ordinal, if α is a limit ordinal we take the direct limit that is the union of $\langle \mathfrak{q}_\beta : \beta < \alpha \rangle$. By Claim 3.3, part (3), it exists and is $\leq_{pr}$-above every $\mathfrak{q}_\beta$ for $\beta < \alpha$. We now choose $\mathfrak{p}_{\alpha,1}$, $\bar{\mathcal{C}} = \langle \underset{\sim}{\mathcal{C}}_{\alpha,\zeta} : \zeta < \zeta(*) \rangle$.

Let us define $\mathfrak{p}_{\alpha,1}$ by choosing $\lambda_{\alpha,\zeta} = |\mathbb{P}_\zeta[\mathfrak{p}_{\alpha,0}]|^{\aleph_0}$ and $\mathbb{P}_{\alpha,\zeta} = \{f : f$ is a finite function from $\lambda_{\alpha,\zeta}$ to $\{0,1\}\}$ and $\mathbb{P}_\zeta[\mathfrak{p}_{\alpha,1}] = \mathbb{P}_\zeta[\mathfrak{p}_{\alpha,0}] \times \mathbb{P}_{\alpha,\zeta}$ and $\underset{\sim}{\eta}^{\mathfrak{p}_{\alpha,0}} = \underset{\sim}{\eta}^{\mathfrak{p}_{\alpha,1}}$. Since Γ_ζ has free amalgamation $\mathfrak{p}_{\alpha,0} \leq_{pr} \mathfrak{p}_{\alpha,1} \in K^*$. Clearly $\Vdash_{\mathbb{P}_0[\mathfrak{p}_{\alpha,1}]} \langle \underset{\sim}{\mathfrak{c}}_\beta, \underset{\sim}{\mathfrak{d}}_\beta : \beta < \alpha \rangle$ is an initial segment of a play of $\Game_\kappa(\mathbf{q})$.

So, by absoluteness, $\Vdash_{\mathbb{P}_0[\mathfrak{p}_{\alpha,1}]} \langle \underset{\sim}{\mathfrak{d}}_\beta : \beta < \alpha \rangle$ is $\leq_{\mathbf{q}}$-linearly ordered. Hence by 1.14 there is a $\mathbb{P}_0[\mathfrak{p}_{\alpha,1}]$-name $\underset{\sim}{\mathcal{C}}_\alpha$ of a $\mathbf{q}$-nice family which (is forced to) include $\{\underset{\sim}{\mathfrak{d}}_\beta : \beta < \alpha\}$.

We now can choose $\underset{\sim}{\mathcal{C}}_{\alpha,\zeta}$ by induction on $\zeta < \zeta(*)$ such that

(a) $\Vdash_{\mathbb{P}_\zeta[\mathfrak{p}_{\alpha,1}]} \underset{\sim}{\mathcal{C}}_{\alpha,\zeta}$ is a $\mathbf{q}$-nice family,

(b) $\underset{\sim}{\mathcal{C}}_{\alpha,0} = \underset{\sim}{\mathcal{C}}_\alpha$ and $\varepsilon < \zeta \rightarrow \Vdash_{\mathbb{P}_\zeta[\mathfrak{p}_{\alpha,1}]} \underset{\sim}{\mathcal{C}}_{\alpha,\varepsilon} \subseteq \underset{\sim}{\mathcal{C}}_{\alpha,\zeta}$,

(c) if $\zeta = \varepsilon + 1$ then $(\mathbb{P}^{\mathfrak{p}_{\alpha,1}}_{\zeta(*)} * \mathbb{A}_{\mathbf{q}}(\underset{\sim}{\mathcal{C}}_{\alpha,\varepsilon}), \mathbb{P}^{\mathfrak{p}_{\alpha,1}}_{\zeta(*)} * \mathbb{A}_{\mathbf{q}}(\underset{\sim}{\mathcal{C}}_{\alpha,\zeta}), \underset{\sim}{\eta}^{\mathfrak{p}_{\alpha,0}}_\zeta) \in \Gamma_\varepsilon$ is above $\mathfrak{p}_{\alpha,1}$.

We carry out the induction by using the Definition 2.9 Part 1 of $\perp$ and thereafter Claim 1.17 to get from directedness to niceness. Or we use Claim 2.10.

Now define $\mathfrak{p}_{\alpha,2} \in K^*$ by choosing $\mathbb{P}_\zeta[\mathfrak{p}_{\alpha,2}] = (\mathbb{P}_\zeta[\mathfrak{p}_{\alpha,1}]) * \mathbb{A}(\underset{\sim}{\mathcal{C}}_{\alpha,\zeta})$. Clearly $\mathfrak{p}_{\alpha,1} \leq_{pr} \mathfrak{p}_{\alpha,2} \in K^*$ so $\mathbb{P}_{\zeta(*)}[\mathfrak{p}_{\alpha,2}] = (\mathbb{P}_{\zeta(*)}[\mathfrak{p}_{\alpha,1}]) * \mathbb{A}(\underset{\sim}{\mathcal{C}}_{\alpha,\zeta(*)})$, where $\underset{\sim}{\mathcal{C}}_{\alpha,\zeta(*)} = \bigcup\{\underset{\sim}{\mathcal{C}}_{\alpha,\zeta} : \zeta < \zeta(*)\}$.

It satisfies the c.c.c by Claim 1.17(1). Lastly let us choose a move $\underset{\sim}{\mathfrak{c}}_\alpha$ for the non-empty player the generic of $\mathbb{A}(\underset{\sim}{\mathcal{C}}_{\alpha,\zeta(*)})$, which is an $\mathbb{A}(\underset{\sim}{\mathcal{C}}_{\alpha,\zeta})$-name for every $\zeta < \zeta(*)$ that is, it is $\bigcup\{\bar{\mathfrak{c}}^p \restriction n^p : p = (n^p, \bar{\mathfrak{c}}^p) \in \underset{\sim}{G}(\mathbb{A}_{\mathbf{q}}(\mathcal{C}_{\alpha,\zeta}))\}$. So half of clause (t_3) holds.

So there is $\mathfrak{p}_{\alpha,3} \in K$ which is by $\leq_{K^*}$ above $\mathfrak{p}_{\alpha,2}$ and forces $\underset{\sim}{St}(\langle \underset{\sim}{\mathfrak{c}}_\beta : \beta \leq \alpha \rangle)$ to be equal to some $\mathbb{P}^{\mathfrak{p}_{\alpha,\zeta}}_\zeta$-name $\underset{\sim}{\bar{\mathfrak{d}}}^\alpha$.

However we cannot in general choose $\mathfrak{q}_\alpha = \mathfrak{p}_{\alpha,3}$, $\underset{\sim}{\bar{\mathfrak{d}}}_\alpha = \underset{\sim}{\bar{\mathfrak{d}}}^\alpha$ as possible $\underset{\sim}{\bar{\mathfrak{d}}}^\alpha$ is not a $\mathbb{P}_0[\mathfrak{p}_{\alpha,3}]$-name. Since $\zeta(*)$ is uncountable, for some $\zeta_\alpha < \zeta(*)$ the name $\underset{\sim}{\bar{\mathfrak{d}}}^\alpha$ is a $\mathbb{P}_{\zeta_\alpha}[\mathfrak{p}_{\alpha,3}]$-name. So we can find $\mathfrak{p}_{\alpha,4}$ such that $\mathfrak{p}_{\alpha,3} \leq_{\mathrm{pr}} \mathfrak{p}_{\alpha,4}$ and $\langle \mathcal{C}^*_{\alpha,\zeta} : \zeta \in [\zeta_\alpha, \zeta(*)) \rangle$ such that for every $\zeta \in [\zeta_\alpha, \zeta(*))$ we have $\mathcal{C}^*_{\alpha,\zeta}$ is a $\mathbb{P}_\zeta[\mathfrak{p}_{\alpha,4}]$-name of a $\mathbf{q}$-nice family to which $\bar{\mathfrak{d}}^\alpha$ is forced to belong increasing with ζ such that

$$(\mathbb{P}_\varepsilon[\mathfrak{p}_{\alpha,4}], \mathbb{P}_{\varepsilon+1}[\mathfrak{p}_{\alpha,4}], \underset{\sim}{\eta}_\varepsilon) \leq_\Gamma (\mathbb{P}_\varepsilon[\mathfrak{p}_{\alpha,4}] * \mathbb{A}_{\mathbf{q}}(\underset{\sim}{\mathcal{C}}^*_{\alpha,\varepsilon}), \mathbb{P}_{\varepsilon+1}[\mathfrak{p}_{\alpha,4}] * \mathbb{A}_{\mathbf{q}}(\underset{\sim}{\mathcal{C}}^*_{\alpha,\varepsilon+1}), \underset{\sim}{\eta}_\varepsilon).$$

Let $\mathfrak{p}_{\alpha,5} \in K^*$ be defined as follows: $\mathbb{P}_\varepsilon[\mathfrak{p}_{\alpha,5}]$ is $\mathbb{P}_\varepsilon[\mathfrak{p}_{\alpha,4}]$ if $\varepsilon < \zeta_\alpha$ and is $\mathbb{P}_\varepsilon[\mathfrak{p}_{\alpha,4}] * \mathbb{A}_{\mathbf{q}}(\underset{\sim}{\mathcal{C}}^*_{\alpha,\varepsilon})$ if $\varepsilon \in [\zeta_\alpha, \zeta(*))$. $\mathcal{C}^*_{\alpha,\zeta(*)} = \bigcup\{\mathcal{C}^*_{\alpha,\zeta} : \zeta < \zeta(*)\}$.

Now clearly $\mathfrak{p}_{\alpha,4} \leq_{\rm pr} \mathfrak{p}_{\alpha,5}$ and let $\underset{\sim}{\bar{\mathfrak{d}}}_\alpha$ be the generic of $\mathbb{A}_{\mathbf{q}}(\underset{\sim}{\mathcal{C}}^*_{\alpha,\zeta(*)})$. Now by Claim 1.20 we have that $\Vdash_{\mathbb{P}_0[\mathfrak{p}_{\alpha,5}]/\mathbb{P}_\varepsilon[\mathfrak{p}_{\alpha,1}]}$ "$\underset{\sim}{\bar{\mathfrak{d}}}_\alpha$ is generic for $\mathbb{A}_{\mathbf{q}}(\underset{\sim}{\mathcal{C}}^*_{\alpha,\zeta(*)})$". Lastly we define $\mathfrak{q}_\alpha$ by

$$\mathbb{P}_\varepsilon[\mathfrak{q}_\alpha] = \begin{cases} (\mathbb{P}_\varepsilon[\mathfrak{p}_{\alpha,5}] * \text{ (the subforcing of} \mathbb{A}_{\mathbf{q}}(\underset{\sim}{\mathcal{C}}^*_{\alpha,\zeta(*)}) & \\ \text{ generated by } \underset{\sim}{\bar{\mathfrak{d}}}_\alpha), & \text{if } \varepsilon \leq \zeta_\alpha, \\ \mathbb{P}_\varepsilon[\mathfrak{p}_{\alpha,5}], & \text{if } \varepsilon > \zeta_\alpha. \end{cases} \tag{3.1}$$

Now in step κ, also $\bar{\mathfrak{c}}_\kappa$ exists and is $\leq_{\mathbb{P}}$, below all the $\bar{\mathfrak{d}}_\alpha$, $\alpha < \kappa$, which is a contradiction to St being a winning strategy for the empty player. □

Theorem 3.9. *1. Assume that* $\mathbf{q} \perp \bar{\Gamma}$, *as in 1.1,* $\bar{\Gamma}$ *be a sequence of Knaster forcing bigness notions* Γ_ζ *with free amalgamation and the* $(< \lambda)$*-LS. as in 2.1,* $K^* = K^*_{\lambda,\zeta(*)}(\bar{\Gamma})$. *Then recalling* $\text{inv}_{\rm gm}(\mathbf{q})$ *from Definition 0.1(4) we have* $\Vdash_{\mathfrak{K}^* * \underset{\sim}{\mathbb{P}}}$ $\text{inv}_{\rm gm}(\mathbf{q}) = \lambda$.

2. If $\bar{\Gamma}$ *has free amalgamation and* $\zeta(*) = \sup\{\zeta < \zeta(*) : \Gamma_\zeta = \Gamma\}$, *then* $\Vdash_{\mathfrak{K}^* * \underset{\sim}{\mathbb{P}}}$ *"for some* $Y \subseteq {}^\omega\omega$ $|Y| = \text{cf}(\zeta(*))$, *for every* $\nu \in {}^\omega\omega$ *(of* $\nu \in {}^\gamma\omega$ *for some* $\gamma < \zeta(*)$*) some* $\eta \in Y$ *is* Γ*-big over* $\{\nu\}$*". For* $\Gamma = \Gamma_{\rm ud}$ *this means* $\eta \not\leq^* \nu$.

Proof. 1. Assume towards a contradiction that we have $\kappa = \text{cf}(\kappa) \in (\aleph_0, \lambda)$ and $\mathfrak{q} \in \mathfrak{K}^*$ such that

$$\mathfrak{q} \Vdash_{\mathfrak{K}^*} p \Vdash_{\underset{\sim}{\mathbb{P}}} \text{inv}_{\rm gm}(\mathbf{q}) = \kappa.$$

So, for some $\underset{\sim}{St}$ we have that $\mathfrak{q} \Vdash_{\mathfrak{K}^*} p \Vdash_{\underset{\sim}{\mathbb{P}}}$ "$\underset{\sim}{St}$ is a name for a winning strategy for the empty player in the game $\Game_\kappa(\mathbf{q})$". But this contradicts the previous claim.

2. is Claim 3.5. □

Equation (3.1) is probably easier to state in terms of complete boolean algebras, because we speak about the generated subforcing. The following remark gives the translation.

Remark 3.10. 1. If $\mathfrak{q} \in K$ and if $\hat{\mathfrak{q}}$ is defined by $\underset{\sim}{\eta}^{\hat{\mathfrak{q}}}_\zeta = \underset{\sim}{\eta}^{\mathfrak{q}}_\zeta$ and $\mathbb{P}^{\hat{\mathfrak{q}}}_\zeta =$ the completion of $\mathbb{P}^{\mathfrak{q}}_\zeta$, then $\mathfrak{q} \leq_{\rm pr} \hat{\mathfrak{q}}$.

2. For a forcing notion $\mathbb{P}$ let its completion $\hat{\mathbb{P}}$ consist of all $\{\mathcal{J} : \mathcal{J} \subseteq \mathbb{P}$ is an antichain$\}$, ordered by $\mathcal{I} \leq \mathcal{J}$ iff $(\forall p \in \mathcal{I})(\exists q \in \mathcal{J})(p \leq_{\mathbb{P}} q)$.

Identifying $\{p\}$ with p we have that $\mathbb{P} \lessdot \hat{\mathbb{P}}$, $\mathbb{P}$ is dense in $\hat{\mathbb{P}}$ and $\hat{\mathbb{P}}$ is a quasi order, there are usually many equivalent elements.

4. Examples

Definition 4.1. *The Hechler forcing in the creature framework.* $\mathbf{q}_H = \mathbf{q} = (K, \Sigma, \mathrm{nor}, \mathrm{val})$ *is given by* $K = \{\mathfrak{c} = (m_{\mathrm{dn}}, m_{\mathrm{up}}, h) : (\exists n < \omega) m_{\mathrm{dn}} = n, m_{\mathrm{up}} = n+1, h \colon \{n\} \to \omega\}$. $\mathrm{dom}(\Sigma) = K$ *and* $\Sigma(m_{\mathrm{dn}}, m_{\mathrm{up}}, h) = \{(m_{\mathrm{dn}}, m_{\mathrm{up}}, g) : g(m_{\mathrm{dn}}) \geq h(m_{\mathrm{dn}})\}$, *and* $\mathrm{nor}(m_{\mathrm{dn}}, m_{\mathrm{up}}, h) = m_{\mathrm{dn}}$, $\mathrm{val}(\mathfrak{c}) = \{(n, h(n))\}$.

Observation 4.2. *1.* $\mathbf{q}_H$ *is a forgetful pcrf as in Definitions 1.1 and 1.2.*
2. $\mathrm{inv}_{\mathrm{cm}}(\mathbf{q}_{\mathrm{H}}) = \mathrm{inv}_{\mathrm{gm}}(\mathbf{q}_{\mathrm{H}}) = \mathfrak{b}$.
3. $\leq_{\mathbf{q}}$ *is the usual* $\leq^*$ *on* ${}^\omega\omega$ *if we identify* $\bar{\mathfrak{c}} = \langle \mathfrak{c}_n : n < \omega \rangle \in \mathbf{Q}$ *with* $f_{\bar{\mathfrak{c}}} \in {}^\omega\omega$, $f_{\bar{\mathfrak{c}}}(m_{\mathrm{dn}}) = h(m_{\mathrm{dn}})$ *if* $(m_{\mathrm{dn}}, m_{\mathrm{up}}, h) \in \mathrm{range}(\bar{\mathfrak{c}})$.
4. $\mathbf{q}_H$ *is not so interesting, as in our framework for* $\mathcal{C}$ *being* $\mathbf{q}$*-nice,* $\mathbb{A}_{\mathbf{q}}(\mathcal{C})$ *will add a dominating real and hence increase* $\mathfrak{b}$.

Definition 4.3. $\mathbf{q}_{\mathrm{gr}} = (K, \Sigma, \mathrm{nor}, \mathrm{val})$ *where*

- *(a)* $K = \{\mathfrak{c} : \mathfrak{c} = (m^{\mathfrak{c}}_{\mathrm{dn}}, m^{\mathfrak{c}}_{\mathrm{up}}, u^{\mathfrak{c}}), m^{\mathfrak{c}}_{\mathrm{dn}} < m^{\mathfrak{c}}_{\mathrm{up}} < \omega, \emptyset \neq u^{\mathfrak{c}} \subseteq [m^{\mathfrak{c}}_{\mathrm{dn}}, m^{\mathfrak{c}}_{\mathrm{up}})\}$
- *(b)* $\mathfrak{c} \in \Sigma(\mathfrak{c}_0, \dots, \mathfrak{c}_{k-1})$ *iff* $\mathfrak{c}_\ell \in K$ *and for* $\ell < k-1$, $m^{\mathfrak{c}_\ell}_{\mathrm{up}} = m^{\mathfrak{c}_{\ell+1}}_{\mathrm{dn}}$, *and* $m^{\mathfrak{c}}_{\mathrm{dn}} = m^{\mathfrak{c}_0}_{\mathrm{dn}}$ *and* $m^{\mathfrak{c}}_{\mathrm{up}} = m^{\mathfrak{c}_{k-1}}_{\mathrm{up}}$ *and for some non-empty* $v \subseteq \{0, \dots, k-1\}$ *we have* $u^{\mathfrak{c}} = \bigcup_{\ell \in v} u^{\mathfrak{c}_\ell}$.
- *(c)* $\mathrm{nor}(\mathfrak{c}) = m^{\mathfrak{c}}_{\mathrm{dn}}$ *for* $\mathfrak{c} \in K$, $\mathrm{val}(\mathfrak{c}) = u_{\mathfrak{c}}$.

Remark: $\mathbf{Q}^{tr}_{\mathbf{q}_{\mathrm{gr}}}$ is called Matet forcing, see [3]. Close to Hindman [7].

Observation 4.4. *1.* $\mathbf{q}_{\mathrm{gr}}$ *is a forgetful pcrf.*
2. $\mathrm{inv}_{\mathrm{gm}}(\mathbf{q}_{\mathrm{gr}}) \leq \mathfrak{g}$.

Proof. 1. Straightforward.
2. Let $\kappa = \mathfrak{g}$. By the definition of $\mathfrak{g}$ we can find $\langle \mathcal{A}_\alpha : \alpha < \kappa \rangle$ such that

- (a) $\mathcal{A}_\alpha \subseteq [\omega]^{\aleph_0}$,
- (b) if $A \subseteq^* B \in \mathcal{A}_\alpha$ and A is infinite, then $A \in \mathcal{A}_\alpha$,
- (c) if $n_0 < n_1 < \dots$ then for some infinite set $w \subseteq \omega$ we have $\bigcup\{[n_i, n_{i+1}) : i \in w\} \in \mathcal{A}_\alpha$,
- (d) for no $B \in [\omega]^{\aleph_0}$ do we have $(\forall \alpha < \kappa)(\exists A \in \mathcal{A}_\alpha)(B \subseteq A)$.

Now we describe a strategy $\mathbf{ST}$ for the empty player in the game $\partial_\kappa(\mathbf{q}_{\mathrm{gr}})$. If the play up to the β-th move has been $\langle (\bar{\mathfrak{c}}_\beta, \bar{\mathfrak{d}}_\beta) : \beta < \alpha \rangle$ and the non-empty player chooses $\bar{\mathfrak{c}}_\alpha$, then the empty player chooses $\bar{\mathfrak{d}}_\alpha$ such that

$$\bar{\mathfrak{c}}_\alpha \leq_{\mathbf{q}_{\mathrm{gr}}} \bar{\mathfrak{d}}_\alpha \wedge \bigcup\{u[\bar{\mathfrak{d}}_{\alpha,n}] : n < \omega\} \in \mathcal{A}_\alpha. \tag{4.1}$$

This is possibe as if we let $m_n = m_{\rm dn}[\bar{\mathfrak{c}}_{\alpha,n}]$ then $m_n < m_{n+1} < \dots$ and $m_0 = 0$ hence for some infinite $w \subseteq \omega$ we have $\bigcup\{[m_i, m_{i+1}) : i \in w\} \in \mathcal{A}_\alpha$ and letting $i_0 < i_1 < i_2 < \dots$ list w, we choose $\mathfrak{d}_{\alpha,\ell} = (m_{i_{\ell-1}+1}, m_{i_\ell+1}), u[\bar{\mathfrak{d}}_{\alpha,\ell}] = u[\bar{\mathfrak{c}}_{\alpha.i_\ell}]$, stipulating $m_{i_{-1}+1} = 0$ so $\langle \mathfrak{d}_{\alpha,\ell} : \ell < \omega\rangle$ is as required.

Suppose that $\langle(\bar{\mathfrak{c}}_\beta, \bar{\mathfrak{d}}_\beta) : \beta < \kappa\rangle$ have been chosen and there is $\bar{\mathfrak{c}}_\kappa = \langle \mathfrak{c}_{\kappa,n} : n < \omega\rangle \in \mathbf{Q}^{tr}_{\mathbf{q}_{\rm gr}}$ such that $\alpha < \kappa \to \bar{\mathfrak{c}}_\alpha \leq_{\mathbf{q}_{\rm gr}} \bar{\mathfrak{c}}_\kappa$. We shall derive a contradiction.

Let the set $B = \bigcup\{u[\mathfrak{c}_{\kappa,n}] : n < \omega\}$. This an infinite subset of ω by the definition of $\mathbf{q}_{\rm gr}$. Also for each $\alpha < \kappa$ clearly $\bar{\mathfrak{c}}_\kappa$ is above $\bar{\mathfrak{c}}_{\alpha+1}$, hence above $\mathfrak{d}_\alpha$, hence $B \subseteq^* \bigcup\{u[\mathfrak{d}_{\alpha,n}] : n < \omega\} \in \mathcal{A}_\alpha$, so $B \in \mathcal{A}_\alpha$. Hence $B \in \bigcup\{\mathcal{A}_\alpha : \alpha < \kappa\}$, contradicting the choice of $\langle \mathcal{A}_\alpha : \alpha < \kappa\rangle$ $\square_{4.4}$

5. ON THE CONSISTENCY OF $\mathfrak{b} \ll \mathfrak{g}$

We now turn to a specific problem.

Theorem 5.1. *Let $\aleph_0 < \kappa = {\rm cf}(\kappa) < \lambda = \lambda^{<\lambda}$. Then for some notion of forcing $\mathfrak{K} * \underset{\sim}{\mathbb{P}}$ of cardinality λ in $\mathbf{V}^{\mathfrak{K} * \underset{\sim}{\mathbb{P}}}$ we have $\mathfrak{b} = \kappa$ and $\mathfrak{g} = \lambda$.*

Definition 5.2. *For $\mathcal{C} \subseteq \mathbf{Q}^{tr}_{\mathbf{q}_{\rm gr}}$ directed and a a filter $D(\mathcal{C}) = \{\{m_{\rm dn}(\mathfrak{c}_n) : n \in \omega\} : \bar{\mathfrak{c}} = \langle \mathfrak{c}_n : n \in \omega\rangle \in \mathcal{C}\}$ we define*

$$\begin{aligned}\mathbf{Q}^{tr}_D = &\{(w, \bar{\mathfrak{c}}) : w \in [\omega]^{<\aleph_0}, \bar{\mathfrak{c}} = \langle \mathfrak{c}_n : n < \omega\rangle \in \mathcal{C}, m_{\rm up}(\mathfrak{c}_n) \leq m_{\rm dn}(\mathfrak{c}_{n+1}),\\ &\sup(w) < m_{\rm dn}(\mathfrak{c}_0), \{m_{\rm dn}(\mathfrak{c})_n : n < \omega\} \in D\}\end{aligned} \tag{5.1}$$

and order like in $\leq^{tr}_{\mathbf{q}}$ in the true ordering.

We use the Definitions from Section 1 for this $\mathbf{q}_{\rm gr}$ and for $\mathbf{Q}^{tr}_{\mathbf{q}_{\rm gr}}$.

Lemma 5.3. $\mathbf{q}_{\rm gr} \perp \Gamma_{\rm ud}$.

Proof. This is a relative of the Main Lemma on page 263 in [5]. We show how to find our scenario in Blass' and Shelah's work: If

(a) $(\mathbb{P}_1, \mathbb{P}_1^+, \underset{\sim}{\eta}) \in \Gamma_{\rm ud}$ and

(b) $\mathbb{P}_1 \lessdot \mathbb{P}_2$ and $\mathbb{P}_2/\mathbb{P}_1$ is ${\rm Cohen}_A$ for some set A and $\mathbb{P}_2 \cap \mathbb{P}_1^+ = \mathbb{P}_1$,

(c) $\underset{\sim}{\mathcal{C}_2}$ is a $\mathbb{P}_2$-name and $\mathbb{P}_2$ forces that it is $\mathbf{q}_{\rm gr}$-nice,

then we can find $(\mathbb{P}_2^+, f, \underset{\sim}{\mathcal{C}_2^+})$ such that

(α) $\mathbb{P}_1^+ \lessdot \mathbb{P}_2^+$,

(β) f is a complete embedding of $\mathbb{P}_2$ into $\mathbb{P}_2^+$ over $\mathbb{P}_1$,

(γ) $\Vdash_{\mathbb{P}_2^+}$ "$f(\underset{\sim}{\mathcal{C}}_2) \subseteq \underset{\sim}{\mathcal{C}}_2{}^+ \wedge \underset{\sim}{\mathcal{C}}_2{}^+$ is $\mathbf{q}_{\mathrm{gr}}$-nice",

(δ) $(\mathbb{P}_1, \mathbb{P}_1^+, \underset{\sim}{\eta}) \leq_{\Gamma_{\mathrm{ud}}} (f(\mathbb{P}_2), \mathbb{P}_2^+, \underset{\sim}{\eta}) \leq_{\Gamma_{\mathrm{ud}}} (f(\mathbb{P}_2) * \mathbb{A}(f(\underset{\sim}{\mathcal{C}}_2)), \mathbb{P}_2^+ * \mathbb{A}(\underset{\sim}{\mathcal{C}}_2{}^+), \underset{\sim}{\eta})$.

Indeed we can choose with a simple witness: $\mathbb{P}_2^+ = (\mathbb{P}_1^+ \times_{\mathbb{P}_1} \mathbb{P}_2) \times \mathrm{Cohen}_\lambda$, $\lambda = (|\mathbb{P}_1^+| + |\mathbb{P}_2|)^{\aleph_0}$, $f = id$.

Since Cohen reals do not add dominating functions, $(\mathbb{P}_1, \mathbb{P}_1^+, \underset{\sim}{\eta}) \leq_{\Gamma_{\mathrm{ud}}} (f(\mathbb{P}_2), \mathbb{P}_2^+, \underset{\sim}{\eta})$. Now for the second part we use the Main Lemma with $M = V^{\mathbb{P}_2}$, $M' = V^{\mathbb{P}_2^+}$. $Q(\mathcal{U})$ has the rôle of $\mathbb{A}_{\mathbf{q}_{\mathrm{gr}}}(\mathcal{C}_2)$ and $Q(\mathcal{U}')$ has the rôle of $\mathbb{A}_{\mathbf{q}_{\mathrm{gr}}}(\mathcal{C}_2^+)$. The $\mathbf{q}_{\mathrm{gr}}$-niceness of $\mathcal{C}$ gives the desired maximality of $\mathcal{C}$ like being an ultrafilter. We first look for $\mathcal{C}^+$ being $\mathbf{q}$-directed. If this were not possible, then finitely many elements would witness this. So this translates into Blass Shelah proof. In the end we can extend the found $\mathbf{q}$-directed $\mathcal{C}^+$ to a $\mathbf{q}$-nice $\mathcal{C}^+$ using Cohen reals as in Conclusion 1.17 Part 1. □

Proof. 5.1: We choose $\bar{\Gamma} = \langle \Gamma_i \,:\, i < \kappa \rangle$ so $\zeta(*) = \kappa$ and $i < \kappa \to \Gamma_i = \Gamma_{\mathrm{ub}}$. Using our present λ we take the forcing notion $\mathfrak{K}^* * \underset{\sim}{\mathbb{P}}$. It does not collapse any cardinals by 3.6 and no cofinality is changed and clearly for some $(\mathfrak{q}, \emptyset) \in \mathfrak{K}^* * \mathbb{P}$ we have $(\mathfrak{q}, \emptyset) \Vdash_{\mathfrak{K}^* * \mathbb{P}} 2^{\aleph_0} = \lambda \wedge MA_{<\kappa}$, so $\mathfrak{b} \geq \kappa$ in the extension. Now by Claim 3.7, for every $(\mathfrak{q}, \emptyset)$, $(\mathfrak{q}, \emptyset) \Vdash_{\mathfrak{K}^* * \mathbb{P}} \{\underset{\sim}{\eta}_i \,:\, i < \kappa\}$ is unbounded, hence $\mathfrak{b} = \kappa$. By Claim 3.9 $\mathrm{inv}_{\mathrm{gm}}(\mathbf{q}_{\mathrm{gr}}) = \lambda$ and by Fact 4.4 Part 2, $\Vdash_{\mathfrak{K}^* * \mathbb{P}} \mathfrak{g} \geq \lambda$, hence since $2^\omega = \lambda$, $\mathfrak{g} = \lambda$. □

REFERENCES

[1] Uri Abraham. Proper forcing. In Matthew Foreman, Akihiro Kanamori, and Menachem Magidor, editors, *Handbook of Set Theory*. Kluwer, To appear.

[2] Tomek Bartoszyński and Haim Judah. *Set Theory, On the Structure of the Real Line.* A K Peters, Wellesley, Massachusetts, 1995.

[3] Andreas Blass. Applications of superperfect forcing and its relatives. In Juris Steprāns and Steve Watson, editors, *Set Theory and its Applications*, volume 1401 of *Lecture Notes in Mathematics*, pages 18–40, 1989.

[4] Andreas Blass and Saharon Shelah. There may be $\mathrm{P}_{\aleph_1}$- and $\mathrm{P}_{\aleph_2}$-points and the Rudin-Keisler ordering may be downward directed. *Ann. Pure Appl. Logic*, 33:213–243, [BsSh:242], 1987.

[5] Andreas Blass and Saharon Shelah. Ultrafilters with small generating sets. *Israel Journal of Mathematics*, 65:259–271, 1989.

[6] Martin Goldstern. Tools for your forcing construction. In Haim Judah, editor, *Set Theory of the Reals*, volume 6 of *Israel Mathematical Conferences Proceedings*, pages 305–360, 1993.

[7] Neil Hindman. Finite sums from sequences within cells of a partition of N. *J. Combin. Theory Ser. A*, 17:1–11, 1974.

[8] Haim Judah and Andrzej Rosłanowski. On Shelah's amalgamation. In Haim Judah, editor, *Set Theory of the Reals*, volume 6 of *Israel Mathematical Conferences Proceedings*, pages 385–415, 1993.

[9] Andrzej Rosłanowski and Saharon Shelah. *Norms on Possibilities I: Forcing with Trees and Creatures*, volume 141 (no. 671) of *Memoirs of the American Mathematical Society*. 1999, [RoSh:470].

[10] Saharon Shelah. Are $\mathfrak{a}$ and $\mathfrak{d}$ your cup of tea? [Sh:700]. *Acta Mathematica*, accepted.

[11] Saharon Shelah. Long iterations for the continuum, [Sh:707]. *Archive for Mathematical Logic*, submitted.

[12] Saharon Shelah. Vive la différence III. *Israel Journal of Mathematics*, submitted, [Sh:509].

[13] Saharon Shelah. On cardinal invariants of the continuum. In *Axiomatic set theory (Boulder, Colo., 1983)*, volume 31 of *Contemp. Mathematics*, pages 183–207, [Sh:207]. Amer. Math. Soc., Providence, RI, 1984. Proceedings of the Conference in Set Theory, Boulder, June 1983; ed. Baumgartner J., Martin, D. and Shelah, S.

[14] Saharon Shelah. *Proper and Improper Forcing, 2nd Edition.* Springer, 1998.

Heike Mildenberger, Universität Wien, Institut für formale Logik, Währinger Str. 25, A-1090 Vienna, Austria

Saharon Shelah, Institute of Mathematics, The Hebrew University of Jerusalem, Givat Ram, 91904 Jerusalem, Israel, and Mathematics Department, Rutgers University, New Brunswick, NJ, USA

E-mail address: `heike@logic.univie.ac.at`

E-mail address: `shelah@math.huji.ac.il`